



***Streptococcus pneumoniae* serotype distribution in low- and middle-income countries of South Asia: Do we need to revisit the pneumococcal vaccine strategy?**

Priya Dhawale, Sanket Shah, Kaushal Sharma, Deepa Sikriwal, Varnik Kumar, Arnabjyoti Bhagawati, Sakshi Dhar, Pratiksha Shetty & Syed Ahmed

To cite this article: Priya Dhawale, Sanket Shah, Kaushal Sharma, Deepa Sikriwal, Varnik Kumar, Arnabjyoti Bhagawati, Sakshi Dhar, Pratiksha Shetty & Syed Ahmed (2025) *Streptococcus pneumoniae* serotype distribution in low- and middle-income countries of South Asia: Do we need to revisit the pneumococcal vaccine strategy?, Human Vaccines & Immunotherapeutics, 21:1, 2461844, DOI: [10.1080/21645515.2025.2461844](https://doi.org/10.1080/21645515.2025.2461844)

To link to this article: <https://doi.org/10.1080/21645515.2025.2461844>



© 2025 The Author(s). Published with license by Taylor & Francis Group, LLC.



Published online: 25 Feb 2025.



Submit your article to this journal [↗](#)



Article views: 702



View related articles [↗](#)



View Crossmark data [↗](#)

REVIEW



Streptococcus pneumoniae serotype distribution in low- and middle-income countries of South Asia: Do we need to revisit the pneumococcal vaccine strategy?

Priya Dhawale^{a*}, Sanket Shah^{b*}, Kaushal Sharma^c, Deepa Sikriwal^d, Varnik Kumar^d, Arnabjyoti Bhagawati^d, Sakshi Dhar^d, Pratiksha Shetty^e, and Syed Ahmed^f

^aGlobal Business Development, Techinvention Lifecare Private Limited, Mumbai, India; ^bStrategic Medical Affairs, Techinvention Lifecare Private Limited, Mumbai, India; ^cStrategic Projects, Techinvention Lifecare Private Limited, Mumbai, India; ^dResearch and Development, Techinvention Lifecare Private Limited, Mumbai, India; ^eRegulatory Affairs, Techinvention Lifecare Private Limited, Mumbai, India; ^fBusiness Development and Strategy, Techinvention Lifecare Private Limited, Mumbai, India

ABSTRACT

S. pneumoniae serotypes responsible for pneumococcal disease differ with respect to disease severity, invasiveness, antimicrobial susceptibility, geographies, immunization history, age groups, and with time. Although PCVs have blunted the pneumococcal disease burden, they are plagued with numerous challenges, especially the emergence of NVTs. In this review, we show that there are diverse serotypes, especially NVTs, responsible for causing pneumococcal diseases in LMICs of South Asia across different studies conducted between 2012 and 2024. We propose that pharmaceutical/biotech companies should tailor/customize the PCVs as per the region-specific serotype prevalence based on surveillance data. Furthermore, protein-based vaccines, or WCVs, have been explored and can serve as viable alternatives to address the limitations associated with PCVs. However, robust studies are warranted in different geographies to demonstrate its efficacy and safety in clinical trials as well as the real-world effectiveness of these promising candidates.

ARTICLE HISTORY

Received 30 August 2024
Revised 20 January 2025
Accepted 30 January 2025

KEYWORDS

Streptococcus pneumoniae; invasive pneumococcal disease; pneumococcal vaccines; antimicrobial resistance; South Asia

Introduction

Streptococcus pneumoniae (*S. pneumoniae*) is a gram-positive bacteria that colonizes the upper respiratory tract of humans.^{1,2} Globally, it represents an important cause of morbidity and mortality in children and adults.³ It is responsible for causing invasive pneumococcal disease (IPD), viz., meningitis, septicemia, bacteremia, and bacteremic pneumonia, as well as non-IPD, such as acute otitis media (AOM) and non-bacteremic pneumonia, in both adults and children.^{4,5} The disease burden (incidence and mortality rates) is highest in children <5 years and adults >65 years, as well as in individuals with compromised immune system or chronic medical conditions.^{5,6} The World Health Organization (WHO) estimates that *S. pneumoniae* is responsible for killing > 300,000 children under 5 years of age globally every year, with the majority of them occurring in developing countries.⁷ It is also the leading cause of bacterial pneumonia globally.⁷ Antibiotics have been used as front-line agents for the treatment of *S. pneumoniae* infections, but the emergence of antibiotic-resistant strains is a major concern for clinicians.^{8,9} The WHO has included *S. pneumoniae* as one of the 12 priority pathogens, exhibiting increased resistance to antibiotics.^{10,11} *S. pneumoniae* produces several virulence factors, and the capsular polysaccharide is considered to be the most important one, enabling binding to the cells and evading the host immune system.^{5,6,12,13} More than 90 serotypes have been identified based on these capsular polysaccharides, and only a subset of them are responsible for causing pneumococcal

disease.^{5,6,14} These serotypes also form the basis of pneumococcal conjugate vaccines (PCVs), and only a fraction of the serotypes are included in the vaccine formulation because of the complexity involved in the manufacturing of PCVs.^{5,15,16} Pneumococcal disease is endemic globally, but the distribution of *S. pneumoniae* serotypes differs with respect to their disease severity, invasiveness, antimicrobial susceptibility, geographies, immunization history of individuals, age groups, and with time.^{4,5,16–19}

Pneumococcal carriage is seen more frequently in children than adults, and thus children are considered to be the main reservoir and transmitter of this pathogen.¹³ In low- and middle-income countries (LMICs), *S. pneumoniae* has three times higher colonization as compared to those in high-income countries (HICs).²⁰ The burden of pneumococcal diseases remains high in LMICs.²¹ Globally, there are over 1,400 cases of pneumonia per 100,000 children annually, with the greatest incidence occurring in South Asia (2,500 cases/100,000 children) and West and Central Africa (1,620 cases/100,000 children).²² South Asian countries have the highest mortality due to IPD.²³ Population-based studies in South Asian countries have shown that *S. pneumoniae* was responsible for 12.8% of invasive bacterial diseases, whereas retrospective hospital-based studies showed that 28% of them were due to *S. pneumoniae*.²³

The implementation of PCVs has played a significant role in reducing the burden of pneumococcal disease globally

CONTACT Syed Ahmed  syed@techinvention.biz  Business Development and Strategy, Techinvention Lifecare Private Limited, #1004, The Summit Business Park, Off WEH Metro Station, Andheri Kurla Road, Andheri East, Mumbai 400093, India.

*These authors contributed equally to this work and share first authorship.

This article has been republished with minor changes. These changes do not impact on the academic content of the article.

© 2025 The Author(s). Published with license by Taylor & Francis Group, LLC.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.

through direct protection of vaccinated individuals and indirect protection of unvaccinated individuals by reducing the nasopharyngeal carriage and transmission of vaccine-type (VT) pneumococci.^{24,25} Vaccines also serve as an effective strategy to address the burgeoning issue of antimicrobial resistance (AMR) in *S. pneumoniae*.⁹ As per the VIEW-hub Report: Global Vaccine Introduction and Implementation, 159 countries have introduced PCVs into their national immunization programs (NIPs) as of August 2024.²⁶ PCVs have also been introduced in many LMICs with the support of Gavi, the Vaccine Alliance.²⁷ Evaluating the impact of vaccines in LMICs is of paramount importance because of differences in pneumococcal epidemiology, including transmission rates.²⁸

Despite the remarkable success achieved with PCVs, the prevalence of IPD has increased by non-vaccine serotypes (NVTs), partially offsetting the total disease reduction.^{4,29–32} This phenomenon, referred to as serotype replacement, is due to the greater serotype diversity observed in developing countries.²⁹ Although this phenomenon was initially considered controversial, it is now well established.³³ These NVTs may replace the VTs in the upper airways, resulting in increased transmission and disease, as well as undermining the effectiveness of the currently available PCVs.^{19,34} They have the tendency to be highly invasive in nature as well as exhibit high levels of resistance to antibiotics.^{29,35} Two mechanisms for serotype replacement have been suggested, which are described in detail by Lo et al. and van Tonder et al.^{30,36} Hence, it becomes imperative to monitor the serotype data post-PCV introduction to identify new emerging serotypes and gain insights into the evolving epidemiology of *S. pneumoniae*.^{4,37} Also, robust vaccine strategies are warranted to address the issue of NVTs.³⁸

In this comprehensive review, we bring forth the distribution and distinctive characteristics of pneumococcal serotypes in LMICs of South Asia. It sheds light on the prevailing trend of NVTs gaining prominence globally, showcasing their distinct features across the region. An elaborate overview of the current pneumococcal vaccine landscape is provided, highlighting the merits and demerits of PCVs, along with alternative strategies (protein-based and whole-cell vaccines [WCVs]) that are being investigated to address the limitations of PCVs. Furthermore, we provide valuable recommendations for future vaccine design and development to mitigate the threat posed by these NVTs.

Methods

We conducted a comprehensive literature search on PubMed to identify relevant studies that were published between 2012 and 2024 (at the time of writing the manuscript) using the keywords, viz., “pneumococcal disease,” “pneumococcal serotypes,” “pneumococcal conjugate vaccines,” “invasive pneumococcal disease,” “non-invasive pneumococcal disease,” and the respective South Asian country name (Afghanistan, Bangladesh, Sri Lanka, Nepal, Bhutan, Pakistan, India, and Maldives). Studies that were conducted between 2012 to 2024 were included in our analysis. Also, studies that commenced before 2012 but concluded in or after 2012 were also included in our analysis. Studies published in native/regional languages other than English were excluded. Table 1 highlights the *S.*

pneumoniae serotypes reported in LMICs of South Asia from different studies conducted between 2012 and 2024 (at the time of writing this manuscript). Table 2 highlights the different characteristics of NVTs observed in this region.

Pneumococcal vaccine landscape

Approved pneumococcal polysaccharide vaccines (PPSVs) and PCVs

There are two types of pneumococcal vaccines available that are clinically available, viz., PPSVs and PCVs, targeting multiple serotypes.^{12,102} PPSVs stimulate the B cells by cross-linking the B-cell receptors (BCR), which leads to the production of immunoglobulins.¹⁰³ This type of immune response is usually a T cell-independent response, which implies that these polysaccharides are capable of activating B cells in the absence of T helper cells.^{104,105} These polysaccharides cannot be processed and presented in association with major histocompatibility complex (MHC) class II molecules and cannot be recognized by T helper cells.¹⁰⁴ Due to the lack of T cell collaboration, the antibodies are of low affinity and are composed mainly of immunoglobulin M (IgM) with limited isotype switching to immunoglobulin G (IgG) and immunoglobulin A (IgA).¹⁰⁴ It results in a lack of production of new memory B cells along with depletion of the memory B cell pool, resulting in hyporesponsiveness to future doses of vaccine.¹⁰³ PPSV23 was licensed in the United States of America (USA) in 1983.⁵ It included 23 serotypes, which were responsible for 85% to 90% of IPD globally.²⁹ It is approved for use in individuals 50 years of age or older and persons aged ≥ 2 years who are at increased risk for pneumococcal disease.¹⁰⁶ PPSV23's protective efficacy has been consistently demonstrated in several studies as well as meta-analysis.^{107–111} In immunocompetent individuals, the vaccine has an efficacy of 65% against pneumococcal infections with serotypes included in it.¹¹² The effectiveness against IPD in immunocompetent adults ranges from 56% to 81% but is lower in individuals with compromised immune status.¹² It reduces the severity of community-acquired pneumonia (CAP) but does not prevent it. It does not reduce the incidence of noninvasive pneumonia.¹² Moreover, it does not generate immunological memory in children <2 years, and hence it is not effective in this population.¹² Also, it does not reduce pneumococcal carriage.^{12,112}

To overcome this drawback and elicit a protective immune response in children <2 years, capsular polysaccharide vaccines conjugated to a protein carrier were developed.^{5,112} Conjugation of the polysaccharide to a suitable carrier protein converts it into a T cell-dependent antigen.¹¹³ This conjugation of the capsular polysaccharide directs the processing of the carrier protein by the polysaccharide-specific B cells, followed by the presentation of the peptides to carrier-peptide specific T cells in association with MHC class II molecules.¹⁰³ When B cells receive help from T cells, they proliferate and differentiate into plasma cells, with class switching to IgG, and memory B cells. These memory B cells undergo rapid proliferation and differentiate into plasma cells upon subsequent encounter with the antigen, producing high antibody titers.^{103,113} A 7-valent PCV (Pneumovax, Pfizer) conjugated to cross-reactive material 197 (CRM₁₉₇) was developed and licensed in the United States (U.S.) and the European Union (EU) in 2000 and 2001, respectively.^{12,102} The vaccine

Table 1. Pneumococcal serotypes reported in LMICs of South Asia from different studies conducted between 2012 to 2024.

Studies included ¹	Age groups ²	Serotypes/Serogroups reported ³	Serotypes covered under PCV13 ⁴	NVTs	Mixed Serotypes	Countries and references	Remarks
- Serotype distribution and AMR patterns of pneumococcal isolates collected from individuals with IPD and non-IPD and from carriage studies - Customized PCR for invasive pneumococcal serotype determination - Comparison of <i>S. pneumoniae</i> serotype distribution, AMR patterns, and carriage from community children and children with clinical pneumonia; health care and non-healthcare workers; and pre- and post-Hajj pilgrims - Carriage of <i>S. pneumoniae</i> serotypes in vaccinated and unvaccinated populations - Evaluating the burden of bacterial meningitis - Distribution and comparison of <i>S. pneumoniae</i> serotypes in healthy and diseased individuals - Comparison of <i>S. pneumoniae</i> serotype distribution in individuals with and without pneumonia - Identification of risk factors associated with nasopharyngeal colonization of <i>S. pneumoniae</i> - Observation and/or comparison of <i>S. pneumoniae</i> serotype distribution, AMR patterns, and genomic changes observed before and after PCV introduction <i>S. pneumoniae</i> carriage, serotype distribution, and AMR patterns in other disease conditions - Identifying pneumococcal serotypes and evaluating the prevalence and risk factors for non-IPD - Effect of antibiotics on invasive and non-invasive pneumococcal isolates - Molecular characterization and serotype distribution of antibiotic-resistant IPD isolates	0 to >65 years	1, 2, 3, 4, 5, 6, 7, 8, 11, 13, 14, 15, 16, 17, 18, 20, 21, 24, 27, 29, 31, 33, 34, 36, 37, 38, 39, 40, 41, 42, 45, 46, 48, 10A, 10B, 10C, 10F, 10G, 10H, 10I, 10J, 10K, 10L, 10M, 10N, 10O, 10P, 10Q, 10R, 10S, 10T, 10U, 10V, 10W, 10X, 10Y, 10Z, 11A, 11B, 11C, 11D, 11E, 11F, 11G, 11H, 11I, 11J, 11K, 11L, 11M, 11N, 11O, 11P, 11Q, 11R, 11S, 11T, 11U, 11V, 11W, 11X, 11Y, 11Z, 12A, 12B, 12C, 12D, 12E, 12F, 12G, 12H, 12I, 12J, 12K, 12L, 12M, 12N, 12O, 12P, 12Q, 12R, 12S, 12T, 12U, 12V, 12W, 12X, 12Y, 12Z, 13A, 13B, 13C, 13D, 13E, 13F, 13G, 13H, 13I, 13J, 13K, 13L, 13M, 13N, 13O, 13P, 13Q, 13R, 13S, 13T, 13U, 13V, 13W, 13X, 13Y, 13Z, 14A, 14B, 14C, 14D, 14E, 14F, 14G, 14H, 14I, 14J, 14K, 14L, 14M, 14N, 14O, 14P, 14Q, 14R, 14S, 14T, 14U, 14V, 14W, 14X, 14Y, 14Z, 15A, 15B, 15C, 15D, 15E, 15F, 15G, 15H, 15I, 15J, 15K, 15L, 15M, 15N, 15O, 15P, 15Q, 15R, 15S, 15T, 15U, 15V, 15W, 15X, 15Y, 15Z, 16A, 16B, 16C, 16D, 16E, 16F, 16G, 16H, 16I, 16J, 16K, 16L, 16M, 16N, 16O, 16P, 16Q, 16R, 16S, 16T, 16U, 16V, 16W, 16X, 16Y, 16Z, 17A, 17B, 17C, 17D, 17E, 17F, 17G, 17H, 17I, 17J, 17K, 17L, 17M, 17N, 17O, 17P, 17Q, 17R, 17S, 17T, 17U, 17V, 17W, 17X, 17Y, 17Z, 18A, 18B, 18C, 18D, 18E, 18F, 18G, 18H, 18I, 18J, 18K, 18L, 18M, 18N, 18O, 18P, 18Q, 18R, 18S, 18T, 18U, 18V, 18W, 18X, 18Y, 18Z, 19A, 19B, 19C, 19D, 19E, 19F, 19G, 19H, 19I, 19J, 19K, 19L, 19M, 19N, 19O, 19P, 19Q, 19R, 19S, 19T, 19U, 19V, 19W, 19X, 19Y, 19Z, 20A, 20B, 20C, 20D, 20E, 20F, 20G, 20H, 20I, 20J, 20K, 20L, 20M, 20N, 20O, 20P, 20Q, 20R, 20S, 20T, 20U, 20V, 20W, 20X, 20Y, 20Z, 21A, 21B, 21C, 21D, 21E, 21F, 21G, 21H, 21I, 21J, 21K, 21L, 21M, 21N, 21O, 21P, 21Q, 21R, 21S, 21T, 21U, 21V, 21W, 21X, 21Y, 21Z, 22A, 22B, 22C, 22D, 22E, 22F, 22G, 22H, 22I, 22J, 22K, 22L, 22M, 22N, 22O, 22P, 22Q, 22R, 22S, 22T, 22U, 22V, 22W, 22X, 22Y, 22Z, 23A, 23B, 23C, 23D, 23E, 23F, 23G, 23H, 23I, 23J, 23K, 23L, 23M, 23N, 23O, 23P, 23Q, 23R, 23S, 23T, 23U, 23V, 23W, 23X, 23Y, 23Z, 24A, 24B, 24C, 24D, 24E, 24F, 24G, 24H, 24I, 24J, 24K, 24L, 24M, 24N, 24O, 24P, 24Q, 24R, 24S, 24T, 24U, 24V, 24W, 24X, 24Y, 24Z, 25A, 25B, 25C, 25D, 25E, 25F, 25G, 25H, 25I, 25J, 25K, 25L, 25M, 25N, 25O, 25P, 25Q, 25R, 25S, 25T, 25U, 25V, 25W, 25X, 25Y, 25Z, 26A, 26B, 26C, 26D, 26E, 26F, 26G, 26H, 26I, 26J, 26K, 26L, 26M, 26N, 26O, 26P, 26Q, 26R, 26S, 26T, 26U, 26V, 26W, 26X, 26Y, 26Z, 27A, 27B, 27C, 27D, 27E, 27F, 27G, 27H, 27I, 27J, 27K, 27L, 27M, 27N, 27O, 27P, 27Q, 27R, 27S, 27T, 27U, 27V, 27W, 27X, 27Y, 27Z, 28A, 28B, 28C, 28D, 28E, 28F, 28G, 28H, 28I, 28J, 28K, 28L, 28M, 28N, 28O, 28P, 28Q, 28R, 28S, 28T, 28U, 28V, 28W, 28X, 28Y, 28Z, 29A, 29B, 29C, 29D, 29E, 29F, 29G, 29H, 29I, 29J, 29K, 29L, 29M, 29N, 29O, 29P, 29Q, 29R, 29S, 29T, 29U, 29V, 29W, 29X, 29Y, 29Z, 30A, 30B, 30C, 30D, 30E, 30F, 30G, 30H, 30I, 30J, 30K, 30L, 30M, 30N, 30O, 30P, 30Q, 30R, 30S, 30T, 30U, 30V, 30W, 30X, 30Y, 30Z, 31A, 31B, 31C, 31D, 31E, 31F, 31G, 31H, 31I, 31J, 31K, 31L, 31M, 31N, 31O, 31P, 31Q, 31R, 31S, 31T, 31U, 31V, 31W, 31X, 31Y, 31Z, 32A, 32B, 32C, 32D, 32E, 32F, 32G, 32H, 32I, 32J, 32K, 32L, 32M, 32N, 32O, 32P, 32Q, 32R, 32S, 32T, 32U, 32V, 32W, 32X, 32Y, 32Z, 33A, 33B, 33C, 33D, 33E, 33F, 33G, 33H, 33I, 33J, 33K, 33L, 33M, 33N, 33O, 33P, 33Q, 33R, 33S, 33T, 33U, 33V, 33W, 33X, 33Y, 33Z, 34A, 34B, 34C, 34D, 34E, 34F, 34G, 34H, 34I, 34J, 34K, 34L, 34M, 34N, 34O, 34P, 34Q, 34R, 34S, 34T, 34U, 34V, 34W, 34X, 34Y, 34Z, 35A, 35B, 35C, 35D, 35E, 35F, 35G, 35H, 35I, 35J, 35K, 35L, 35M, 35N, 35O, 35P, 35Q, 35R, 35S, 35T, 35U, 35V, 35W, 35X, 35Y, 35Z, 36A, 36B, 36C, 36D, 36E, 36F, 36G, 36H, 36I, 36J, 36K, 36L, 36M, 36N, 36O, 36P, 36Q, 36R, 36S, 36T, 36U, 36V, 36W, 36X, 36Y, 36Z, 37A, 37B, 37C, 37D, 37E, 37F, 37G, 37H, 37I, 37J, 37K, 37L, 37M, 37N, 37O, 37P, 37Q, 37R, 37S, 37T, 37U, 37V, 37W, 37X, 37Y, 37Z, 38A, 38B, 38C, 38D, 38E, 38F, 38G, 38H, 38I, 38J, 38K, 38L, 38M, 38N, 38O, 38P, 38Q, 38R, 38S, 38T, 38U, 38V, 38W, 38X, 38Y, 38Z, 39A, 39B, 39C, 39D, 39E, 39F, 39G, 39H, 39I, 39J, 39K, 39L, 39M, 39N, 39O, 39P, 39Q, 39R, 39S, 39T, 39U, 39V, 39W, 39X, 39Y, 39Z, 40A, 40B, 40C, 40D, 40E, 40F, 40G, 40H, 40I, 40J, 40K, 40L, 40M, 40N, 40O, 40P, 40Q, 40R, 40S, 40T, 40U, 40V, 40W, 40X, 40Y, 40Z, 41A, 41B, 41C, 41D, 41E, 41F, 41G, 41H, 41I, 41J, 41K, 41L, 41M, 41N, 41O, 41P, 41Q, 41R, 41S, 41T, 41U, 41V, 41W, 41X, 41Y, 41Z, 42A, 42B, 42C, 42D, 42E, 42F, 42G, 42H, 42I, 42J, 42K, 42L, 42M, 42N, 42O, 42P, 42Q, 42R, 42S, 42T, 42U, 42V, 42W, 42X, 42Y, 42Z, 43A, 43B, 43C, 43D, 43E, 43F, 43G, 43H, 43I, 43J, 43K, 43L, 43M, 43N, 43O, 43P, 43Q, 43R, 43S, 43T, 43U, 43V, 43W, 43X, 43Y, 43Z, 44A, 44B, 44C, 44D, 44E, 44F, 44G, 44H, 44I, 44J, 44K, 44L, 44M, 44N, 44O, 44P, 44Q, 44R, 44S, 44T, 44U, 44V, 44W, 44X, 44Y, 44Z, 45A, 45B, 45C, 45D, 45E, 45F, 45G, 45H, 45I, 45J, 45K, 45L, 45M, 45N, 45O, 45P, 45Q, 45R, 45S, 45T, 45U, 45V, 45W, 45X, 45Y, 45Z, 46A, 46B, 46C, 46D, 46E, 46F, 46G, 46H, 46I, 46J, 46K, 46L, 46M, 46N, 46O, 46P, 46Q, 46R, 46S, 46T, 46U, 46V, 46W, 46X, 46Y, 46Z, 47A, 47B, 47C, 47D, 47E, 47F, 47G, 47H, 47I, 47J, 47K, 47L, 47M, 47N, 47O, 47P, 47Q, 47R, 47S, 47T, 47U, 47V, 47W, 47X, 47Y, 47Z, 48A, 48B, 48C, 48D, 48E, 48F, 48G, 48H, 48I, 48J, 48K, 48L, 48M, 48N, 48O, 48P, 48Q, 48R, 48S, 48T, 48U, 48V, 48W, 48X, 48Y, 48Z, 49A, 49B, 49C, 49D, 49E, 49F, 49G, 49H, 49I, 49J, 49K, 49L, 49M, 49N, 49O, 49P, 49Q, 49R, 49S, 49T, 49U, 49V, 49W, 49X, 49Y, 49Z, 50A, 50B, 50C, 50D, 50E, 50F, 50G, 50H, 50I, 50J, 50K, 50L, 50M, 50N, 50O, 50P, 50Q, 50R, 50S, 50T, 50U, 50V, 50W, 50X, 50Y, 50Z, 51A, 51B, 51C, 51D, 51E, 51F, 51G, 51H, 51I, 51J, 51K, 51L, 51M, 51N, 51O, 51P, 51Q, 51R, 51S, 51T, 51U, 51V, 51W, 51X, 51Y, 51Z, 52A, 52B, 52C, 52D, 52E, 52F, 52G, 52H, 52I, 52J, 52K, 52L, 52M, 52N, 52O, 52P, 52Q, 52R, 52S, 52T, 52U, 52V, 52W, 52X, 52Y, 52Z, 53A, 53B, 53C, 53D, 53E, 53F, 53G, 53H, 53I, 53J, 53K, 53L, 53M, 53N, 53O, 53P, 53Q, 53R, 53S, 53T, 53U, 53V, 53W, 53X, 53Y, 53Z, 54A, 54B, 54C, 54D, 54E, 54F, 54G, 54H, 54I, 54J, 54K, 54L, 54M, 54N, 54O, 54P, 54Q, 54R, 54S, 54T, 54U, 54V, 54W, 54X, 54Y, 54Z, 55A, 55B, 55C, 55D, 55E, 55F, 55G, 55H, 55I, 55J, 55K, 55L, 55M, 55N, 55O, 55P, 55Q, 55R, 55S, 55T, 55U, 55V, 55W, 55X, 55Y, 55Z, 56A, 56B, 56C, 56D, 56E, 56F, 56G, 56H, 56I, 56J, 56K, 56L, 56M, 56N, 56O, 56P, 56Q, 56R, 56S, 56T, 56U, 56V, 56W, 56X, 56Y, 56Z, 57A, 57B, 57C, 57D, 57E, 57F, 57G, 57H, 57I, 57J, 57K, 57L, 57M, 57N, 57O, 57P, 57Q, 57R, 57S, 57T, 57U, 57V, 57W, 57X, 57Y, 57Z, 58A, 58B, 58C, 58D, 58E, 58F, 58G, 58H, 58I, 58J, 58K, 58L, 58M, 58N, 58O, 58P, 58Q, 58R, 58S, 58T, 58U, 58V, 58W, 58X, 58Y, 58Z, 59A, 59B, 59C, 59D, 59E, 59F, 59G, 59H, 59I, 59J, 59K, 59L, 59M, 59N, 59O, 59P, 59Q, 59R, 59S, 59T, 59U, 59V, 59W, 59X, 59Y, 59Z, 60A, 60B, 60C, 60D, 60E, 60F, 60G, 60H, 60I, 60J, 60K, 60L, 60M, 60N, 60O, 60P, 60Q, 60R, 60S, 60T, 60U, 60V, 60W, 60X, 60Y, 60Z, 61A, 61B, 61C, 61D, 61E, 61F, 61G, 61H, 61I, 61J, 61K, 61L, 61M, 61N, 61O, 61P, 61Q, 61R, 61S, 61T, 61U, 61V, 61W, 61X, 61Y, 61Z, 62A, 62B, 62C, 62D, 62E, 62F, 62G, 62H, 62I, 62J, 62K, 62L, 62M, 62N, 62O, 62P, 62Q, 62R, 62S, 62T, 62U, 62V, 62W, 62X, 62Y, 62Z, 63A, 63B, 63C, 63D, 63E, 63F, 63G, 63H, 63I, 63J, 63K, 63L, 63M, 63N, 63O, 63P, 63Q, 63R, 63S, 63T, 63U, 63V, 63W, 63X, 63Y, 63Z, 64A, 64B, 64C, 64D, 64E, 64F, 64G, 64H, 64I, 64J, 64K, 64L, 64M, 64N, 64O, 64P, 64Q, 64R, 64S, 64T, 64U, 64V, 64W, 64X, 64Y, 64Z, 65A, 65B, 65C, 65D, 65E, 65F, 65G, 65H, 65I, 65J, 65K, 65L, 65M, 65N, 65O, 65P, 65Q, 65R, 65S, 65T, 65U, 65V, 65W, 65X, 65Y, 65Z, 66A, 66B, 66C, 66D, 66E, 66F, 66G, 66H, 66I, 66J, 66K, 66L, 66M, 66N, 66O, 66P, 66Q, 66R, 66S, 66T, 66U, 66V, 66W, 66X, 66Y, 66Z, 67A, 67B, 67C, 67D, 67E, 67F, 67G, 67H, 67I, 67J, 67K, 67L, 67M, 67N, 67O, 67P, 67Q, 67R, 67S, 67T, 67U, 67V, 67W, 67X, 67Y, 67Z, 68A, 68B, 68C, 68D, 68E, 68F, 68G, 68H, 68I, 68J, 68K, 68L, 68M, 68N, 68O, 68P, 68Q, 68R, 68S, 68T, 68U, 68V, 68W, 68X, 68Y, 68Z, 69A, 69B, 69C, 69D, 69E, 69F, 69G, 69H, 69I, 69J, 69K, 69L, 69M, 69N, 69O, 69P, 69Q, 69R, 69S, 69T, 69U, 69V, 69W, 69X, 69Y, 69Z, 70A, 70B, 70C, 70D, 70E, 70F, 70G, 70H, 70I, 70J, 70K, 70L, 70M, 70N, 70O, 70P, 70Q, 70R, 70S, 70T, 70U, 70V, 70W, 70X, 70Y, 70Z, 71A, 71B, 71C, 71D, 71E, 71F, 71G, 71H, 71I, 71J, 71K, 71L, 71M, 71N, 71O, 71P, 71Q, 71R, 71S, 71T, 71U, 71V, 71W, 71X, 71Y, 71Z, 72A, 72B, 72C, 72D, 72E, 72F, 72G, 72H, 72I, 72J, 72K, 72L, 72M, 72N, 72O, 72P, 72Q, 72R, 72S, 72T, 72U, 72V, 72W, 72X, 72Y, 72Z, 73A, 73B, 73C, 73D, 73E, 73F, 73G, 73H, 73I, 73J, 73K, 73L, 73M, 73N, 73O, 73P, 73Q, 73R, 73S, 73T, 73U, 73V, 73W, 73X, 73Y, 73Z, 74A, 74B, 74C, 74D, 74E, 74F, 74G, 74H, 74I, 74J, 74K, 74L, 74M, 74N, 74O, 74P, 74Q, 74R, 74S, 74T, 74U, 74V, 74W, 74X, 74Y, 74Z, 75A, 75B, 75C, 75D, 75E, 75F, 75G, 75H, 75I, 75J, 75K, 75L, 75M, 75N, 75O, 75P, 75Q, 75R, 75S, 75T, 75U, 75V, 75W, 75X, 75Y, 75Z, 76A, 76B, 76C, 76D, 76E, 76F, 76G, 76H, 76I, 76J, 76K, 76L, 76M, 76N, 76O, 76P, 76Q, 76R, 76S, 76T, 76U, 76V, 76W, 76X, 76Y, 76Z, 77A, 77B, 77C, 77D, 77E, 77F, 77G, 77H, 77I, 77J, 77K, 77L, 77M, 77N, 77O, 77P, 77Q, 77R, 77S, 77T, 77U, 77V, 77W, 77X, 77Y, 77Z, 78A, 78B, 78C, 78D, 78E, 78F, 78G, 78H, 78I, 78J, 78K, 78L, 78M, 78N, 78O, 78P, 78Q, 78R, 78S, 78T, 78U, 78V, 78W, 78X, 78Y, 78Z, 79A, 79B, 79C, 79D, 79E, 79F, 79G, 79H, 79I, 79J, 79K, 79L, 79M, 79N, 79O, 79P, 79Q, 79R, 79S, 79T, 79U, 79V, 79W, 79X, 79Y, 79Z, 80A, 80B, 80C, 80D, 80E, 80F, 80G, 80H, 80I, 80J, 80K, 80L, 80M, 80N, 80O, 80P, 80Q, 80R, 80S, 80T, 80U, 80V, 80W, 80X, 80Y, 80Z, 81A, 81B, 81C, 81D, 81E, 81F, 81G, 81H, 81I, 81J, 81K, 81L, 81M, 81N, 81O, 81P, 81Q, 81R, 81S, 81T, 81U, 81V, 81W, 81X, 81Y, 81Z, 82A, 82B, 82C, 82D, 82E, 82F, 82G, 82H, 82I, 82J, 82K, 82L, 82M, 82N, 82O, 82P, 82Q, 82R, 82S, 82T, 82U, 82V, 82W, 82X, 82Y, 82Z, 83A, 83B, 83C, 83D, 83E, 83F, 83G, 83H, 83I, 83J, 83K, 83L, 83M, 83N, 83O, 83P, 83Q, 83R, 83S, 83T, 83U, 83V, 83W, 83X, 83Y, 83Z, 84A, 84B, 84C, 84D, 84E, 84F, 84G, 84H, 84I, 84J, 84K, 84L, 84M, 84N, 84O, 84P, 84Q, 84R, 84S, 84T, 84U, 84V, 84W, 84X, 84Y, 84Z, 85A, 85B, 85C, 85D, 85E, 85F, 85G, 85H, 85I, 85J, 85K, 85L, 85M, 85N, 85O, 85P, 85Q, 85R, 85S, 85T, 85U, 85V, 85W, 85X, 85Y, 85Z, 86A, 86B, 86C, 86D, 86E, 86F, 86G, 86H, 86I, 86J, 86K, 86L, 86M, 86N, 86O, 86P, 86Q, 86R, 86S, 86T, 86U, 86V, 86W, 86X, 86Y, 86Z, 87A, 87B, 87C, 87D, 87E, 87F, 87G, 87H, 87I, 87J, 87K, 87L, 87M, 87N, 87O, 87P, 87Q, 87R, 87S, 87T, 87U, 87V, 87W, 87X, 87Y, 87Z, 88A, 88B, 88C, 88D, 88E, 88F, 88G, 88H, 88I, 88J, 88K, 88L, 88M, 88N, 88O, 88P, 88Q, 88R, 88S, 88T, 88U, 88V, 88W, 88X, 88Y, 88Z, 89A, 89B, 89C, 89D, 89E, 89F, 89G, 89H, 89I, 89J, 89K, 89L, 89M, 89N, 89O, 89P, 89Q, 89R, 89S, 89T, 89U, 89V, 89W, 89X, 89Y, 89Z, 90A, 90B, 90C, 90D, 90E, 90F, 90G, 90H, 90I, 90J, 90K, 90L, 90M, 90N, 90O, 90P, 90Q, 90R, 90S, 90T, 90U, 90V, 90W, 90X, 90Y, 90Z, 91A, 91B, 91C, 91D, 91E, 91F, 91G, 91H, 91I, 91J, 91K, 91L, 91M, 91N, 91O, 91P, 91Q, 91R, 91S, 91T, 91U, 91V, 91W, 91X, 91Y, 91Z, 92A, 92B, 92C, 92D, 92E, 92F, 92G, 92H, 92I, 92J, 92K, 92L, 92M, 92N, 92O, 92P, 92Q, 92R, 92S, 92T, 92U, 92V, 92W, 92X, 92Y, 92Z, 93A, 93B, 93C, 93D, 93E, 93F, 93G, 93H, 93I, 93J, 93K, 93L, 93M, 93N, 93O, 93P, 93Q, 93R, 93S, 93T, 93U, 93V, 93W, 93X, 93Y, 93Z, 94A, 94B, 94C, 94D, 94E, 94F, 94G, 94H, 94I, 94J, 94K, 94L, 94M, 94N, 94O, 94P, 94Q, 94R, 94S, 94T, 94U, 94V, 94W, 94X, 94Y, 94Z, 95A, 95B, 95C, 95D, 95E, 95F, 95G, 95H, 95I, 95J, 95K, 95L, 95M, 95N, 95O, 95P, 95Q, 95R, 95S, 95T, 95U, 95V, 95W, 95X, 95Y, 95Z, 96A, 96B, 96C, 96D, 96E, 96F, 96G, 96H, 96I, 96J, 96K, 96L, 96M, 96N, 96O, 96P, 96Q, 96R, 96S, 96T, 96U, 96V, 96W, 96X, 96Y, 96Z, 97A, 97B, 97C, 97D, 97E, 97F, 97G, 97H, 97I, 97J, 97K, 97L, 97M, 97N, 97O, 97P, 97Q, 97R, 97S, 97T, 97U, 97V, 97W, 97X, 97Y, 97Z, 98A, 98B, 98C, 98D, 98E, 98F, 98G, 98H, 98I, 98J, 98K, 98L, 98M, 98N, 98O, 98P, 98Q, 98R, 98S, 98T, 98U, 98V, 98W, 98X, 98Y, 98Z, 99A, 99B, 99C, 99D, 99E, 99F, 99G, 99H, 99I, 99J, 99					

Table 2. Characteristics of NVTs observed in LMICs of South Asia from different studies conducted between 2012 and 2024.

NVTs	Characteristics					References
	IPD	Non-IPD	Drug resistance ¹	NP carriage	Unspecified (IPD/Non-IPD) ²	
2	✓	✓	✓	✓	✓	17,43–45,49,50,52,58,63,64,67,68,70,72,76,78,80,84,100
6				✓	✓	48
7	✓	✓				86
8	✓	✓	✓	✓	✓	41–46,49–52,57–59,61–67,69–75,77,84,88,89,92,93,99–101
11*		✓			✓	39,86
13	✓	✓	✓	✓	✓	17,43–47,50–54,56,58–63,67,69–71,73–75,77,80,83–85,89,92,94,100
15	✓					49
16	✓			✓		49,71
17	✓					59,69,77
18*	✓			✓	✓	39,42,65,92
20	✓	✓	✓	✓	✓	17,39,41,43–47,50–53,58,61–63,65,67,69–71,74,75,77,80,83,84,89,90,92,94
21	✓		✓	✓	✓	43,44,46,50–52,54,56,58,61–64,67,68,70–73,83,92
24	✓		✓	✓	✓	41,44,52,53,61,69,70,84
27	✓			✓	✓	17,43,46,50,64,66,71,72,74,75,84
29	✓	✓	✓	✓	✓	17,43,45,46,50–52,61,66,69–72,74,80,90,94,100
31	✓	✓	✓	✓	✓	17,43,44,46,50–52,56,58,62–65,67,70–72,75,78,84–86,89,92,94,100,101
33	✓					49,74
34	✓	✓	✓	✓	✓	39,41,43–47,50–54,56,58,60–65,67,70–75,80,83–85,89,93,94,100,101
36	✓		✓	✓	✓	46,50–52,64,70,72,80,84,90,94,100
37	✓			✓	✓	46,50–52,70,71,75
38	✓	✓	✓	✓	✓	17,43,44,46,50–52,61,64,65,69–75,80,84,89,94,100
39	✓	✓	✓	✓	✓	44–46,50–53,61,66,67,69–71,74,80,83,89,100
40	✓		✓	✓	✓	46,50,51,69,72,77,84
41	✓					52
42				✓	✓	46,50,52,80
45	✓	✓	✓	✓	✓	42,43,45,51,61,64,70,72,75,80,84,100
46	✓					67
48	✓			✓	✓	39,46,49–52,70–72,84,90,94
10A	✓	✓	✓	✓	✓	39,43–46,50–54,56–77,80,83,84,86,89,92–94,100,101
10B	✓		✓	✓	✓	46,50,52,61,70,71,93
10C	✓		✓	✓	✓	17,46,69,70,74,75,83,89,93,100
10F*	✓	✓	✓	✓	✓	43–46,50–52,61,64,66,69–74,77,83,84,92
11A	✓	✓	✓	✓	✓	17,43–46,50–52,61,66,69–75,77,78,80,83–85,89,92–94,101
11B				✓		50,51
11C	✓		✓	✓		46,75
11D	✓			✓		51,69,77,80
11F	✓			✓		43,80
12A	✓				✓	42,43,49,50,52,69,73,80,83
12B	✓			✓		46,70,75
12F	✓	✓	✓	✓	✓	39,41,43,44,46,50–52,57,61,64,69,70,72,74,77,83–85,92,101
15A	✓	✓	✓	✓	✓	43–46,49–52,61,64,68–75,77,80,83,84,89,90,94,100
15B	✓	✓	✓	✓	✓	39,41–43,45,46,50–52,64,66,69–75,77,78,80,83,84,89,90,92,93,99,101
15C*	✓	✓	✓	✓	✓	41,45,46,50–52,64,66,71–74,80,83,84,89,90,94
15F	✓			✓	✓	43,46,50,52,61,64,69,70,72,77,78,80,92
16A				✓		46,50
16C				✓		52
16F	✓	✓	✓	✓	✓	43–46,50–54,56,58,60–65,67,70–75,80,83,84,86,89,90,92,93,101
17A	✓			✓		46,50–52,69,73,77,90,92
17F*	✓	✓	✓	✓	✓	17,39,42–44,46,47,50–52,54,56,58,61–67,69–75,80,84,85,89,90,92,93,99–101
18A	✓	✓	✓	✓	✓	41–43,46,50–52,61,64,66,69,70,72,78,80,83,84,88,100
18B	✓	✓	✓	✓		45,46,50,57,64,69–72,84
18F	✓			✓	✓	43,46,50,61,64,69,70,72,77,93
19B*	✓	✓	✓	✓	✓	45,46,50–52,61,64,66,68,70–72,74,75,84,90,93
19C	✓			✓		49,51,70
19V				✓		97
22A	✓		✓	✓	✓	46,50–52,61,66,70,71,74,75,80,90,100
22F	✓	✓	✓	✓	✓	41,43,44,46,50–52,61,65,69,71–73,75,77,80,84,89,90,92,101
23A*	✓		✓	✓	✓	39,43,44,46,50–53,56,58,61–64,66,67,70–75,80,84,89,90,92,100
23B	✓	✓	✓	✓	✓	44,46,50–54,56–63,67,69–71,75,80,83,85,89,90,93
24A	✓		✓	✓		50–52,64,72,89
24B	✓		✓	✓	✓	17,49,51,52,66,69,74,75,77,80,86,89,90,92
24F	✓		✓	✓	✓	17,42,43,46,50,52,64,66,71–74,89,92–94,100
25A	✓			✓	✓	50,64,69,72,80,84
25F	✓		✓	✓	✓	43,51,52,61,70,78,84,100
28A	✓	✓	✓	✓	✓	43,46,50,51,61,69,70,73,84,85
28F	✓	✓	✓	✓	✓	45,46,50–52,64,70,72,75,80,84
32A				✓	✓	17,52
32F				✓		46,51
33A	✓			✓	✓	46,51,64,72,73,80,92,93
33B	✓		✓	✓	✓	43,46,50–52,61,64,69–73,77,80,84–86,93
33C*	✓		✓	✓	✓	41,43,50–52,66,69–71,74,75,96
33D	✓			✓	✓	41,46,50,52,80

(Continued)

Table 2. (Continued).

NVTs	Characteristics					References
	IPD	Non-IPD	Drug resistance ¹	NP carriage	Unspecified (IPD/Non-IPD) ²	
33F	✓	✓	✓	✓	✓	43,44,46,49–52,64–66,69–72,74,77,80,84,100
35A	✓		✓	✓	✓	42,44,46,50–52,61,64,70–72,75,80,83,84,89,93,100
35B*	✓	✓	✓	✓	✓	17,39,41–47,50–54,56,58–65,67,69–73,75,77,80,83,84,86,89,92,93,101
35C		✓		✓	✓	17,46,50–52,73,80,85
35D	✓		✓			89
35F	✓	✓	✓	✓	✓	39,43,44,46,50–52,61,64,68,70,72,73,80,85,90
41F	✓					72
47F	✓					17,66,74
6C	✓	✓	✓	✓	✓	41,43–47,50–53,56–58,61–64,66,69–75,80,82,84,89,94,100
6D	✓	✓	✓	✓	✓	41,43,45–47,50–52,56,58,61–63,80,84
6F				✓		46
7A	✓			✓		50,69,70,74,75
7B	✓	✓	✓	✓	✓	41–43,45,46,50–52,61,70–72,84,89
7C*	✓		✓	✓	✓	44,46,50–52,61,64,65,70,72,84,89,96
9A	✓			✓	✓	46,64,71–73,80,93
9B	✓		✓			70,75
9L	✓	✓	✓	✓		45,46,50–52,61,84
9N	✓	✓	✓	✓	✓	17,50–52,61,66,69–71,74,77,80,88–90,101

¹Drug resistance refers to resistance to one or more classes of antibiotics.

²Unspecified includes cases that have not mentioned whether the pneumococcal infection was invasive or non-invasive in nature.

³Serogroups/serotypes marked with *in South Asia were responsible for causing deaths.^{39,68,74,96}

⁴Mixed serotypes, as observed in Table 1, have not been included in the analysis because of the inconsistency in how they are named and interpreted across the referenced research articles. This inconsistency limits the ability to accurately characterize mixed serotypes within the study's framework, thus warranting their exclusion to maintain clarity and analytical appropriateness.

Abbreviations: NVTs: Non-vaccine serotypes; IPD: Invasive pneumococcal disease; Non-IPD: Non-invasive pneumococcal disease; NP: Nasopharyngeal.

included serotypes, which were identified as causes of IPD in the U.S.¹³ However, after the introduction of PCV7, a change in the epidemiology of pneumococcal diseases was observed, and NVTs (not included in PCV7) emerged as the most prevalent ones, a phenomenon known as ‘serotype replacement.’^{5,12} For example, serotypes 1 and 5, which were not included in PCV7, were responsible for a high burden of illness and deaths in low-income settings.¹³ Another example was 19A, a problematic serotype, which exhibited antibiotic resistance.^{12,13} This serotype replacement led to the development of PCV10 (Synflorix, GlaxoSmithKline [GSK]) and PCV13 (Pneumovax 13, Pfizer), which were introduced in 2009.^{4,12} Surveillance data from different countries implementing PCVs in their NIPs has prevented disease not only in vaccinated individuals but also in unvaccinated adults and children through herd immunity.¹³ Although PCVs were originally developed for pediatric use, clinical studies conducted in adults have shown their ability to prevent diseases in individuals of all ages through direct protection.¹³ A systematic review and meta-analysis conducted to evaluate the impact of PCV10 and PCV13 in reducing the incidence of CAP in children <5 years demonstrated that these vaccines showed a reduction of 17% and 31% in the hospitalization rates for clinically and radiologically confirmed pneumonia, respectively, in children <24 months.¹¹⁴ In children aged 24–59 months, these vaccines showed a reduction of 9% and 24% in the hospitalization rates for clinically and radiologically confirmed pneumonia, respectively.¹¹⁴ A systematic review evaluating the impact of PCV10 and PCV13 on all-cause, radiologically confirmed, and severe pneumonia hospitalization rates and pneumonia mortality in children 0–9 years old across several WHO regions demonstrated reductions ranging from 7% to 60% for all-cause pneumonia hospitalization, 8% to 90% for severe pneumonia, 12% to 79% for radiologically confirmed

pneumonia, and 45% to 85% for pneumococcal confirmed pneumonia.¹¹⁵ Pneumonia-related mortality saw a reduction of 10% to 78% across different regions.¹¹⁵ A study conducted by Bennett et al. evaluated the impact of PCV10/PCV13 on IPD incidence in children <5 years and adults ≥65 years compared to the pre-PCV period.¹¹⁶ It evaluated 275,582 cases from 47 surveillance sites in 29 countries (primarily HICs [80%]). The study demonstrated that IPD declined by 58% to 79% in children <5 years and by 4% to 29% in adults.¹¹⁶ The Serum Institute of India (SII) has developed an affordable 10-valent PCV (Pneumosil), which provides protection against serotypes that are responsible for causing serious illness in developing countries.¹² Walvax Biotechnology Co., Ltd. (Walvax) has developed and manufactured a 13-valent PCV (Weuphoria), which has been marketed in China, Morocco, and Thailand.¹¹⁷

However, after the introduction of PCV10 and PCV13, the issue of serotype replacement continued, and newer vaccines were required to prevent pneumococcal diseases caused by serotypes 8, 12F, and 22F.^{5,118,119} A systematic review of the serotype distribution of *S. pneumoniae* causing IPD in children in countries that have introduced PCV13 revealed that non-PCV13 serotypes contributed to 42.2% of the childhood IPD cases in the post-PCV era.³⁷ There were differences as per geography, viz., 57.8% in North America, 71.9% in Europe, 45.9% in the Western Pacific, 28.5% in Latin America, and 42.7% in South Africa. The predominant non-PCV13 serotypes were 22F, 12F, 33F, 24F, 15C, 15B, 23B, 10A, and 38.³⁷

In light of this grim situation, two new PCVs, viz., PCV15 (VAXNEUVANCE, Merck) and PCV20 (Pneumovax 20, Pfizer), have been approved by the United States Food and Drug Administration (USFDA) in 2021.^{5,12,120,121} PCV15 is indicated for active immunization for the prevention of invasive disease in individuals 6 weeks of age and older.¹²² PCV20 is

approved for the prevention of invasive diseases in individuals 6 weeks of age and older; prevention of otitis media in individuals 6 weeks through 5 years of age; and active immunization for the prevention of pneumonia in individuals 18 years of age and older.¹²³ Prevnar 20, which was built using Prevnar 13, includes seven additional serotypes that have been associated with antibiotic resistance, heightened disease severity, invasive potential, high case fatality rates, and prevalence in pediatric pneumococcal cases.^{124,125} Moreover, these seven serotypes are among the most common responsible for IPD in the pediatric population in the USA.^{124,125} Biological E Ltd. has developed a 14-valent PCV, viz., PNEUBEVAX 14, which received approval from the Indian drug regulator, Drug Controller General of India (DCGI), in 2022 for manufacturing and commercialization in India.¹²⁶ It offers protection against two new serotypes, 22F and 33F.¹²⁶ A 15-valent PCV has been developed by Tergene Biotech Pvt. Ltd. and manufactured at AuroVaccines Pvt. Ltd., which is a wholly owned subsidiary of Aurobindo Pharma Ltd.¹²⁷ The Indian drug regulator has granted permission to Tergene in 2022 to manufacture and market the vaccine.¹²⁷ In 2024, the USFDA approved CAPVAXIVE (V116), a 21-valent PCV developed by Merck.¹²⁸ The vaccine is indicated for active immunization for the prevention of invasive disease and pneumonia in individuals 18 years of age and older.¹²⁸ It includes eight unique serotypes responsible for approximately 27% of IPD cases in adults 50 years of age and older and approximately 30% in adults 65 years of age and older, as per the Centre for Disease Control and Prevention (CDC) data.¹²⁸

Investigational PPSVs and PCVs

Panacea Biotech has developed an 11-valent PCV, viz., NuCoVac-11, which is currently in phase 3 trial. The company has also developed a PPSV23, which is in preclinical development.¹²⁹ Techinvention Lifecare Pvt. Ltd. has developed a 16-valent PCV that is currently in preclinical studies.¹³⁰ GBP410 is a 21-valent PCV that is jointly developed by SK Bioscience and Sanofi. The vaccine has demonstrated immunogenicity and safety in phase II trials and is expected to enter phase 3 trials in H1 of 2024.¹³¹ AFX3772, a 24-valent PCV developed by Affinivax (acquired by GSK), is currently in phase 2 development.¹³² This vaccine candidate is based on the highly innovative multiple-antigen presenting system (MAPS) platform technology.¹³² A 30-plus-valent pneumococcal candidate vaccine is also in preclinical development.¹³² VAX-24, developed by Vaxcyte, is a 24-valent, broad-spectrum, carrier-sparing PCV that will be entering phase 3.^{133,134} VAX-31, another vaccine developed by Vaxcyte, is a 31-valent PCV, which is expected to enter phase 1/2 clinical study during the fourth quarter of 2023.¹³⁵ VAX-31 and VAX-24 are designed to improve upon the standard-of-care PCVs for both children and adults by providing coverage for serotypes that are responsible for a significant portion of IPD as well as high case-fatality rates, antibiotic resistance, and meningitis.¹³⁴ Preclinical immunogenicity evaluation of a 24-valent PCV developed by Merck has been conducted in adult monkeys.¹³⁶ Inventprise Inc. has developed a 25-valent PCV that is

currently in phase 2 trial.¹³⁷ Table 3 highlights the status of PCVs in LMICs of South Asia. Table 4 gives a list of investigational and approved PPSVs and PCVs, along with their composition as well as their clinical/commercial status.

In light of recent findings, it has become increasingly evident that a decline in immunogenicity against shared serotypes has been noted with the adoption of higher-valency PCVs.¹⁵⁵ This observed trend, while pointing toward a potential decrease in immune response, poses uncertainties regarding its clinical implications.¹⁵⁵ While reduced immunogenicity has been observed among higher- vs. lower-valency PCVs, the trend is inconsistent.¹⁵⁵ Different vaccine components, platforms, technologies, and manufacturing processes can affect serotype-specific immunogenicity.¹⁵⁵ Consequently, further exploration through additional research is needed to fully comprehend the impact and effectiveness of these new vaccines.¹⁵⁵

Exploring various avenues to enhance immunogenicity and overall vaccine efficacy is key in this scenario.¹⁵⁵ Innovative conjugation processes, diverse carriers, strategic use of adjuvants, and advancements in vaccine platforms represent promising approaches that could potentially offset any reduction in immunogenicity.¹⁵⁵ Some of these strategies include choosing appropriate carrier proteins to prevent carrier-induced suppression, such as utilizing single or multiple conserved *S. pneumoniae* proteins or employing enhanced carrier protein (eCRM) to facilitate conjugation reactions outside the primary T cell epitope region.¹⁵⁵ Techniques like preserving the structural integrity of capsular polysaccharides through polysaccharide activation or using specific linker moieties between polysaccharides and proteins are also being explored.¹⁵⁵ Novel vaccine platforms, like MAPS, in vivo bioconjugation, virus-like particle (VLP) technology, and messenger ribonucleic acid (mRNA) technology are being investigated to enhance overall B cell and T cell responses while generating enhanced serotype-specific IgG levels without the need for adjuvants.¹⁵⁵ By striving to maintain or elevate immunogenicity levels and subsequently boost vaccine effectiveness, it's possible to extend the coverage and protection offered by PCVs, especially as their valency increases.¹⁵⁵

VAX-24 contains eCRM, a proprietary carrier protein with non-native amino acids (para-azidomethyl-L-phenylalanine) that undergo site-specific conjugation to pneumococcal polysaccharides activated with a small-molecule linker (dibenzocyclooctyne).¹⁵⁶ Site-specific conjugation utilizing click chemistry allows consistent exposure of T cell epitopes, reduction in carrier protein to pneumococcal polysaccharide ratio, and augments manufacturing process consistency to improve PCVs by increasing the coverage of serotypes and minimizing carrier suppression.¹⁵⁶ VAX-24 has demonstrated promising results in a phase 1/2 trial by targeting a wide array of pneumococcal serotypes without compromising the immunogenicity.¹⁵⁷

Despite the widespread use of PCVs in vaccination programs and continuous advancements in serotype coverage, the underlying reasons for decreased antibody responses, particularly in infants and the elderly, remain incompletely understood.¹⁵⁸ Deeper insights into the molecular and cellular mechanisms governing protective antibody responses against *S. pneumoniae* could potentially inform the development of adjuvants that enhance these responses.¹⁵⁸

Table 3. PCV status in LMICs of South Asia.

Country	Introduced in NIP	Date of introduction	Type of PCV formulation	Dosing schedule ¹	References
Afghanistan	Yes	7 th December 2013	Prenar 13	3+0	138
Bangladesh	Yes	21 st March 2015	Synflorix	3+0	
Sri Lanka	No	NA	NA	NA	
Nepal	Yes	19 th January 2015	Synflorix	2+1	
Bhutan	Yes	1 st January 2019	Prenar 13	2+1	
Pakistan	Yes	9 th October 2012	Prenar 13	3+0	
India	Yes	13 th May 2017	Pneumosil	2+1	

¹3+0: Three primary doses without a booster; 2+1: Two primary doses with one booster.¹³⁹

Abbreviations: NIP: National immunization program; PCV: Pneumococcal conjugate vaccine.

Table 4. List of investigational and approved PPSVs and PCVs.

Sr. No.	Vaccine/ Brand name	Company	Carrier protein	Serotype composition	Clinical/ Commercial status	WHO prequalification	Reference
1	Pneumovax	Merck	–	1, 2, 3, 4, 5, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20, 22F, 23F, and 33F	Approved commercially	–	140
2	PPSV23	Sinovac Biotech Ltd.	–	1, 2, 3, 4, 5, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20, 22F, 23F, and 33F	Approved commercially	–	140
3	PPSV23	Walvax Biotechnology Co. Ltd.	–	1, 2, 3, 4, 5, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20, 22F, 23F, and 33F	Approved commercially	–	12,141
4	Prenar	Pfizer	CRM ₁₉₇	4, 6B, 9V, 14, 18C, 19F, and 23F	Approved commercially	–	12,140
5	Synflorix	GSK	Protein D for serotypes 1, 5, 6B, 7F, 9V, 14, 23F, and 4. TT for serotype 18C and DT for serotype 19.	1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F, and 23F	Approved commercially	Yes	12,140,142
6	Pneumosil	SII	CRM ₁₉₇	1, 5, 6A, 6B, 7F, 9V, 14, 19A, 19F, and 23F	Approved commercially	Yes	12,140,142
7	Prenar 13	Pfizer	CRM ₁₉₇	1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, and 23F	Approved commercially	Yes	12,140,142
8	Weuphoria	Walvax Biotechnology Co., Ltd.	TT	1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, and 23F	Approved commercially	–	12, 117, 140
9	PNEUBEVAX 14	Biological E Ltd.	CRM ₁₉₇	1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19A, 19F, 22F, 23F, and 33F	Approved commercially	–	126,143
10	VAXNEUVANCE	Merck	CRM ₁₉₇	1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 22F, 23F, and 33F	Approved commercially		12,140
11	PCV15	Tergene Biotech Pvt. Ltd. and AuroVaccines Pvt. Ltd.	CRM ₁₉₇	1, 2, 3, 4, 5, 6A, 6B, 7F, 9V, 12F, 14, 18C, 19A, 19F, and 23F	Permission granted by Indian drug regulator to manufacture and market		127,144
12	Prenar 20	Pfizer	CRM ₁₉₇	1, 3, 4, 5, 6A, 6B, 7F, 8, 9V, 10A, 11A, 12F, 14, 15B, 18C, 19A, 19F, 22F, 23F, and 33F	Approved commercially		12,140
13	CAPVAXIVE	Merck	CRM ₁₉₇	3, 6A, 7F, 8, 9N, 10A, 11A, 12F, 15A, 15C, 16F, 17F, 19A, 20, 22F, 23A, 23B, 24F, 31, 33F, and 35B	Approved commercially		128,145
14	NuCoVac-11	Panacea Biotec	CRM ₁₉₇	1, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, and 23F	Phase 3		129,146
15	NuVac-23	Panacea Biotec	–	–	Preclinical		129
16	PneumoWin	Technvention Lifecare Pvt. Ltd.	rCRM ₁₉₇	1, 3, 4, 5, 6A, 6B, 7F, 9V, 12F, 14, 15A, 18C, 19A, 19F, 22F, and 23F	Preclinical		130,147
17	GBP410	SK Bioscience and Sanofi	Serotypes 1, 3 and 5 conjugated to TT. Serotypes 4, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 18C, 19A, 19F, 22F, 23F, and 33F conjugated to CRM ₁₉₇	1, 3, 4, 5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 18C, 19A, 19F, 22F, 23F, and 33F	Completed phase 2		131,148
18	AFX3772	GSK	Rhizavidin-fused carrier proteins	1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F, 2, 8, 9N, 10A, 11A, 12F, 15B, 17F, 20B, 22F, and 33F	Currently in phase 2 development		132,140,149

(Continued)

Table 4. (Continued).

Sr. No.	Vaccine/ Brand name	Company	Carrier protein	Serotype composition	Clinical/ Commercial status	WHO prequalification	Reference
19	VAX-24	Vaxcyte	eCRM	1, 2, 3, 4, 5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20B, 22F, 23F, and 33F	Will be entering phase 3		133,134,140,150
20	PCV24	Merck	CRM ₁₉₇	1, 2, 3, 4, 5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20, 22F, 23F, and 33F	Preclinical development		136
21	PCV25	Inventprise Inc.	CRM ₁₉₇	1, 2, 3, 4, 5, 6B, 6C, 7F, 8, 9N, 9V, 10A, 12F, 14, 15A, 15B, 16F, 18C, 19A, 19F, 22F, 23F, 24F, 33F, and 35B	Phase 2		151,152
22	VAX-31	Vaxcyte	eCRM	1, 2, 3, 4, 5, 6A, 6B, 7C, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15A, 15B, 16F, 17F, 18C, 19A, 19F, 20B, 22F, 23A, 23B, 23F, 31, 33F, and 35B	Expected to enter phase 1/2 clinical trial		135,153,154
23	30-plus valent PCV	GSK	-	-	Preclinical development		132

Abbreviations: WHO: World Health Organization; PPSV: Pneumococcal polysaccharide vaccine; CRM₁₉₇: Cross-reactive material 197; GSK: GlaxoSmithKline; TT: Tetanus toxoid; DT: Diphtheria toxoid; SI: Serum Institute of India; rCRM₁₉₇: Recombinant cross-reactive material 197; eCRM: Enhanced carrier protein.

Moreover, in alignment with the guidelines set forth by the WHO, the evaluation of noninferiority between higher- and lower-valency PCVs relies on specific criteria related to IgG response rates and IgG geometric mean concentrations (GMCs).¹⁵⁹ Each serotype is individually assessed to determine noninferiority, with the WHO underscoring that meeting the predefined criteria for either IgG response rates or GMCs should suffice for regulatory approval.¹⁵⁹ The correlate of protection for IPD is typically measured by noting the levels of vaccine-induced, serotype-specific IgG, with a protective threshold set at 0.35 µg/ml.¹⁵⁹ In 2008, the WHO reviewed data on PCVs, including effectiveness studies in Canada, the United Kingdom (UK), and the USA.¹⁵⁹ The data indicated variability in IgG concentrations and thresholds, with some suggesting alternative thresholds and the correlation of opsonophagocytic activity (OPA) titers for a few serotypes for vaccine effectiveness.¹⁵⁹

Drawbacks

Administration of additional pneumococcal vaccines may be required in adults due to waning immunity. For example, the effectiveness of PPSV23 wanes over several years, and high-risk individuals may require revaccination every 5–10 years.¹¹¹ As mentioned earlier, serotype replacement has been observed with the implementation of PCVs, where niches occupied earlier by VTs have now been colonized by NVTs.^{29,38,160,161} Both PPSVs and PCVs are serotype-based vaccines and therefore elicit a serotype-specific immune response.¹⁶² Geographical differences with respect to serotype distribution have led to reduced vaccine effectiveness when they were implemented in geographies where the serotypes were not covered by the vaccines.²⁹ PCVs have weak efficacy at mucosal sites, such as for pneumonia and otitis media.^{12,38} *S. pneumoniae* undergoes phase variation, wherein the capsule shifts stochastically from high to low expression levels.^{12,38} This is a result of anatomical-site-specific selective pressures in the host environment.^{12,38} The pathogens present on mucosal surfaces are less encapsulated as compared to those present in the blood, thereby exposing surface adhesins.^{12,38} Therefore, the antibodies directed at the capsule are not efficient in

opsonophagocytosis at the mucosal level.^{12,38} Serotype coverage by PCVs in developed countries is substantially higher compared to developing countries, where pneumococcal diseases are caused by a wider spectrum of serotypes.¹⁶² Vaccine failures have been reported following immunization with PCV10 and PCV13.^{162,163} A systematic review published by Oligbu et al. identified serotypes 19B, 6B, and 4 associated with vaccine failure in countries with established PCV programs.¹⁶⁴ PCVs cannot offer protection against the emergence of *S. pneumoniae* strains, which are found among non-typeable isolates.¹⁶⁵ Capsular switching has been observed post the introduction of PCVs.²⁹ Although PCVs provide protection from infection, but not all serotypes are covered with equal potency.¹² For some serotypes, viz., 1, 3, and 5, the levels of functional antibodies against capsular polysaccharides induced by the vaccines and detected by opsonophagocytic killing are not ideal.¹² PCV manufacturing is an extremely complex and expensive process, making it unaffordable for the developing countries, which can lead to low vaccination coverage.^{102,162} One of the glaring examples is the sub-Saharan countries, which still have the highest mortality rates in children <5 years, even after 10 years post-PCV licensure.¹⁰² Increasing the valency of PCV also poses challenges with respect to conjugation.¹⁰² A crucial issue of PCV is the binding of the carrier protein to the capsular polysaccharide to elicit a robust immune response as compared to the capsular polysaccharide alone.¹⁰² Adding additional serotypes to the formulation results in blunting of the immune response to individual serotypes due to carrier suppression.¹² An increase in valency as well as the dosage of the carrier protein in the formulation can lead to immune response interference, both in the vaccine itself and other vaccines that are administered previously, jointly, or subsequently.^{102,166} These interferences result in an imbalance in several arms of the immune response and hence require readjustment in each new vaccine formulation.¹⁰²

Alternative approaches

In light of the above-mentioned drawbacks, novel vaccines based on a non-serotype-independent approach targeting

Table 5. Properties of different *S. pneumoniae* proteins investigated as vaccine candidates.

Protein		Properties	References
Ply		(1) Highly conserved	9,29,170,171
		(2) Immune evasion	
		(3) Triggers inflammation	
		(4) Cell death	
CBPs	PspA	(1) Highly conserved	29,165,172
		(2) Immune evasion	
		(3) Pneumococcal adhesion	
		(4) Binds lactoferrin for critical iron uptake	
	PspC	(1) Highly immunogenic	9,29,173
		(2) Highly conserved	
		(3) Immune evasion	
		(4) Pneumococcal adhesion	
	PcpA	(1) Pneumococcal adhesion	165,174
		(2) Biofilm formation	
	LytA	(1) Immunogenic	165,175
		(2) Highly conserved	
LytB	(3) Facilitates the release of cytotoxin Ply	165	
	(4) Inhibits complement		
	(1) Biofilm formation		
	(2) Attachment to host epithelial cells		
CbpG	(3) Immune evasion	165	
	(4) Well-conserved		
	(1) Pneumococcal adhesion		
	(2) Secreted form can cleave host ECM		
PsaA	CbpM	(1) Binds fibronectin	165
		(1) Oxidative stress resistance	
		(2) Bacterial adhesion	9,176
		(3) Manganese and zinc acquisition	
PHTs	PhtA	(4) Highly conserved	177,178
		(1) Highly conserved	
		(2) Immune system evasion	
		(3) Bacterial dissemination in the lungs	
	PhtB	(1) Conserved protein	179
		(2) Immunogenic	
	PhtD	(1) Highly conserved	165,179–183
		(2) Adherence to nasopharyngeal and epithelial cell lines	
		(3) Inhibits complement	
		(4) Zinc acquisition	
	PhtE	(1) Highly conserved protein	165,179,180
		(2) Adherence to nasopharyngeal and epithelial cell lines	
(3) Inhibits complement			
(1) Iron uptake			
PiuA and PiaA	(2) Highly conserved	9	
	(3) Highly immunogenic		
	(1) Biofilm formation		
PsrP	(2) Lung adhesion and infection	165	
	(1) Highly conserved protein		
Ef-Tu	(2) Enhances pathogenicity through complement inhibition and degradation of ECM	184	

Abbreviations: Ply: Pneumolysin; CBPs: Choline-binding proteins; PspA: Pneumococcal surface protein A; PspC: Pneumococcal surface protein C; PcpA: Pneumococcal choline binding protein A; LytA: N-acetylmuramoyl-L-alanine amidase; LytB: N-acetylglucosaminidase; CbpG: Choline-binding protein G; ECM: Extracellular matrix; CbpM: Choline-binding protein M; PsaA: Pneumococcal surface antigen A; PHTs: Pneumococcal histidine triad proteins; PhtA: Pneumococcal histidine triad protein A; PhtB: Pneumococcal histidine triad protein B; PhtD: Pneumococcal histidine triad protein D; PhtE: Pneumococcal histidine triad protein E; PiuA: Pneumococcal iron uptake A; PiaA: Pneumococcal iron acquisition A; PsrP: Pneumococcal serine-rich repeat protein; Ef-Tu: Elongation factor Tu.

conserved antigens are warranted and have been extensively investigated.^{10,38,102,112,140,162,165} An ideal pneumococcal vaccine with broader coverage, the ability to induce mucosal and systemic immunity, and the ability to prevent primary intranasal colonization and invasive disease is highly desirable.¹² Several protein-based vaccines and whole-cell (live-attenuated and inactivated) vaccines are being evaluated in preclinical and clinical studies.¹⁰ Protein vaccines targeting the highly conserved virulence proteins can confer broad serotype coverage.²⁹ Different types of protein vaccines that are investigated include recombinant proteins (individual or in combination), fusion or hybrid proteins, nanoparticle-packaged fusion or hybrid proteins, recombinant protein-boosted PCVs, and capsular polysaccharide-conjugated recombinant protein carrier.^{162,167} Protein-based vaccines offer several advantages, such as the production

of opsonophagocytic antibodies, providing respiratory tract protection, preventing invasion of the pathogen in different organs, neutralizing pneumolysin cytotoxicity, cross-protection against additional childhood pathogens, and blunting complications arising out of disseminated bacterial infections.^{38,162} It can also serve as a cheaper option for children in developing countries, provide coverage for a wide range of serotypes responsible for IPD, and offer an alternative option to prevent recurrent infections.¹⁴⁰ With respect to manufacturing, there are less hindrances associated with structural differences and variations, a drawback that is commonly observed with conjugate vaccines.¹⁴⁰ On the flip side, protein antigens have numerous limitations. They contain non-repeating conformational epitopes for antibody binding as compared to capsular polysaccharides that have repeating and conformationally less-restricted antigenic

Table 6. Results of protein-based vaccines investigated in animal models.

Protein		Results	References
Ply		● Intramuscular immunization with recombinant PlyD1 protected mice against lethal intranasal challenge with pneumococci and lung injury mediated through Ply challenge. Protection was associated with PlyD1-specific IgG titers and <i>in vitro</i> neutralization titers.	185
		● CaP-ΔA146PLY ¹ nanoparticles induced efficient humoral and cellular immune responses in mice that protected them from both pneumonia and sepsis.	186
		● Intramuscular immunization with dPly protected mice against lethal intranasal challenge with Ply. Also, intranasal immunization in mice inhibited nasopharyngeal colonization after intranasal challenge with homologous or heterologous pneumococcal strain.	187
		● Mice immunized subcutaneously with ΔA146Ply-SP0148 ² elicited high levels of anti-ΔA146Ply and anti-SP0148 IgG antibodies. Mice had a high survival rate when challenged with a lethal dose of <i>S. pneumoniae</i> .	188
		● Subcutaneous immunization of mice with ΔA146 Ply ³ generated high titers of anti-Ply IgGs and protected them from intraperitoneal challenge with TIGR4 <i>S. pneumoniae</i> .	189
		● Intramuscular immunization of mice with PLY-D ⁴ conferred significant protection post-intranasal challenge with different <i>S. pneumoniae</i> serotypes and strains.	190
		● Vaccination with a monovalent formulation of recombinant PlyD1 led to bacterial reductions in both infant and adult mice after a lethal-dose intranasal challenge with serotype 6A.	191
		● Intramuscular immunization of mice with recombinant CbpG provided significant protection against colonization and substantial protection against systemic infection by wild-type TIGR4 cells.	192
		● Subcutaneous immunization with recombinant CbpG protected mice against intraperitoneal challenge with serotype 19F pneumococcal strain.	193
		● Subcutaneous immunization with recombinant CbpM protected mice against intraperitoneal challenge with serotype 19F pneumococcal strain.	193
CBPs	CbpG	● Purified recombinant LytA administered intranasally elicited IgG and IgA antibodies in mice. These mice also showed significantly higher survival rates and lower bacterial carriage in response to <i>S. pneumoniae</i> infection.	194
		● Intraperitoneal immunization with LytB enhanced <i>S. pneumoniae</i> recognition by complement components C1q and C3b, demonstrating that anti-LytB antibodies activated the classical pathway. Vaccination with LytB protected mice against pneumococcal sepsis and invasive pneumonia caused by clinical isolates of serotypes 3 or 23F.	195
	CbpM	● Intranasal immunization with PspC using cholera toxin subunit B as an adjuvant protected mice against intranasal challenge with pneumolysin mutant of D39 or serotype 6B strain BG7322.	196
		● Intraperitoneal immunization of PspC protected mice from intraperitoneal challenge with virulent capsular type 2 pneumococcal strain D39.	197
	LytA	● Intranasal immunization with PspC induced high levels of anti-PspC antibodies in mice. The hyperimmune sera significantly inhibited the adhesion of <i>S. pneumoniae</i> to A549 cells.	198
		● Nasal immunization with <i>L. casei</i> expressing PspC primed the immune system of mice that prompted faster immune responses, resulting in a decrease in pneumococcal colonization.	199
	LytB	● Subcutaneous immunization with purified PspC protected mice against sepsis. The protection was mediated by antibodies cross-reactive with PspA.	200
		● Subcutaneous or intranasal immunization of PspC vaccines did not protect mice against an invasive intranasal challenge with pneumococcal strain ATCC 6303.	201
	PspC/ CbpA	● Vaccination with a monovalent formulation of recombinant PcpA led to bacterial reductions in both infant and adult mice after a lethal-dose intranasal challenge with serotype 6A.	191
		● Antibodies generated post-immunization with recombinant family 1 PspA in humans in a phase 1 trial offered passive protection in mice against a pneumococcal challenge from serotypes 3, 6A, or 6B.	202
	PcpA	● Transcutaneous immunization with PspA + cholera toxin B induced antigen-specific mucosal and systemic immune responses in mice. It significantly reduced the CFU numbers in nasal washes and passages when they were challenged with <i>S. pneumoniae</i> strain EF3030 via the nasal route.	203
		● Intranasal immunization with a nanogel-based PspA nasal vaccine provided protective immunity in mice against intranasal lethal challenge with <i>S. pneumoniae</i> Xen10. It reduced the bacterial colonization and invasion and induced high levels of PspA-specific IgG and nasal and bronchial IgA responses.	204
	PspA	● Mucosal immunization with various nanoparticle PspA formulations protected mice against lethal intranasal challenge <i>S. pneumoniae</i> strain ATCC6303.	205
		● Intranasal immunization with PspA delivered by BLPs enhanced PspA-specific antibody responses in mice and provided significant protection against intranasal challenge with pneumococci of two different clades of both homologous and heterologous PspA families.	206
		● Intranasal immunization with PspA-BLP vaccine ⁵ conferred protection in mice against fatal intranasal challenge with both PspA family 1 and 2 pneumococcal strains regardless of serotype.	207
		● Intramuscular immunization with polymer particles entrapping PspA elicited robust IgG responses both in mice and in rats. Antigen load in microparticles correlated with the antibody titers.	208
		● Mucosal immunization with NP/NCMP PspA4Pro ⁶ provided partial protection against intranasal lethal pneumococcal challenge with a serotype 3 strain expressing PspA from clade 5.	209
		● Intranasal vaccination with PspA and cationic liposomes conferred protective immunity against lethal <i>S. pneumoniae</i> inhalation and improved the survival rate of infected mice. It also induced the production of PspA-specific IgA and IgG by both mucosal and systemic compartments as well as elicited PspA-specific Th17 responses.	210

(Continued)

Table 6. (Continued).

Protein		Results	References
PsaA		Nasal vaccination of macaques with cCHP-trivalent PspA generated PspA-specific IgGs that protected from pneumococcal intratracheal challenge through inhibition of lung inflammation and a reduction in the numbers of bacteria in the lungs.	215
		Subcutaneous immunization with PspA3+2⁸ provided significant protection against intranasal challenge by five pneumococcal strains with different clades in mice.	216
		Nasal immunization of mice with a combination of PspA5 and wP⁹ or wP_{low}¹⁰ conferred protection against intranasal respiratory lethal challenge with <i>S. pneumoniae</i> ATCC6303 or A66.1 strain. Both PspA5-wP and PspA5-wP_{low} vaccines induced high levels of systemic and mucosal antibodies against PspA5.	217
		Subcutaneous immunization of mice with PsaA using either complete Freund's or TiterMax adjuvants significantly protected against challenge from type 3 pneumococcal strain WU2.	218
		Intranasal or oral administration of full-length PsaA in mice demonstrated significant reduction in colonization of nasopharyngeal tissues after intranasal challenge with <i>S. pneumoniae</i> strains compared to controls.	219
		Intraperitoneal immunization with PsaA admixed with BlyS exhibited only modest elevations in PsaA-specific responses. Mice sera obtained post-immunization exhibited high titers of IgG1, IgG2a, IgG2b, and IgG3, but no IgA.	220
		Intranasal immunization with chitosan-PsaA¹¹ generated mucosal and systemic immune responses in mice, and fewer pneumococci were recovered from the immunized mice following intranasal challenge with ATCC 6303 (serotype 3).	221
		Intranasal immunization with chitosan-PsaA generated mucosal and systemic immune responses in mice. Immunized mice had increased protection against AOM following middle ear challenge with pneumococcus serotype 14 and improved survival following intraperitoneal challenge with pneumococcus serotype 3 or serotype 14.	222
PHTs	PhtA	Subcutaneous immunization with full-length PhtA generated antibodies that protected mice against lethal sepsis challenge with serotype 6A or 6B.	178
		Immunization with either N-terminal or C-terminal portion of PhtA also protected mice against serotype 6B challenge.	
	PhtB	- N-terminal half of PhtA induced a protective response against serotype 6A challenge. - Subcutaneous immunization with full-length PhtB generated high levels of antibody titers and protected mice against challenge with serotype 6B.	178
	PhtD	Subcutaneous immunization with recombinant PhtD induced antibodies that protected mice against sepsis challenge with serotype 6A and 6B and highly virulent capsular type 4 strain or with the capsular type 3 strain WU2.	178
		Subcutaneous immunization of PhtD and its C-terminal fragment provided protection against pneumococcal sepsis in mice.	223
		Mice immunized intraperitoneally with PhtD and its C-terminal with OMVs and alum as adjuvants survived after intraperitoneal challenge with <i>S. pneumoniae</i> ATCC6303.	224
		Vaccination with a monovalent formulation of recombinant PhtD led to bacterial reductions in both infant and adult mice after a lethal-dose intranasal challenge with serotype 6A.	191
		Intranasal administration of PhtD in mice conferred protection against pneumococcal colonization through robust serum antibodies and CD4 Th1-biased immune memory.	225
	PhtE	Intranasal administration of PhtE in mice conferred protection against pneumococcal colonization through robust serum antibodies and CD4 Th1-biased immune memory.	225
		Subcutaneous immunization of mice with recombinant PhtE elicits protective immunity against experimental sepsis and pneumonia.	226
PsrP		Antibodies directed against SRR1 and BR regions of recombinant PsrP neutralized bacterial adhesion to lung cells <i>in vitro</i>.	227
Ef-Tu		Passive immunization of mice with PsrP antiserum conferred protection against pneumococcal challenge.	
	- Subcutaneous immunization with recombinant EF-Tu induced the production of inflammatory cytokines and IgG1 and IgG2a antibodies in mice. - Significantly and non-specifically protected mice against lethal intraperitoneal challenge with <i>S. pneumoniae</i> serotypes 2 and 15A.	184	
Pneumococcal proteins explored as carrier proteins			
PLD ¹²		Mice immunized subcutaneously with 19F polysaccharide conjugated with PLD as a carrier protein elicited high antibody titers against the polysaccharide and the carrier protein. Immunized mice effectively cleared the bacteria and had a significantly better survival rate when they were challenged intraperitoneally with type 19F pneumococci.	228
PLD		Subcutaneous immunization of mice with both monovalent and tetravalent conjugate formulations (serotypes 6B, 14, 19F, and 23F) conjugated to PLD as a carrier protein generated high levels of polysaccharide-specific IgGs that demonstrated opsonophagocytic activity for serotypes tested, viz., 6B, 14, 19F, and 23F.	229
Ply		Intraperitoneal immunization of 18C polysaccharide conjugated to recombinant Ply conferred significant protection in mice challenged with type 18C pneumococci.	230
PsaA		Intraperitoneal immunization with Hib polysaccharide vaccine conjugated with recombinant PsaA protein carrier in mice generated anti-PsaA and anti-Hib immune responses. It was able to offer effective protection against AOM caused by pneumococcus.	231
Proteins explored in combination			
PhtD-dPly		Intranasal immunization with PhtD-dPly vaccine protected rhesus macaques from mortality induced by <i>S. pneumoniae</i> challenge.	232
PlyD6 ¹³ , PspA, PsaA, PiuA, RrgB, RrgA, Hsp 60, and Hsp 70		Intramuscular vaccination of rabbits with a multiantigen vaccine comprising PlyD6, PspA, PsaA, PiuA, RrgB, RrgA, Hsp 60, and Hsp 70 demonstrated the killing of vaccine strain TIGR4 in an opsonophagocytic killing assay and heterologous strains 6B, 19F, and 15B.	233
Ply, PspC, and PspA		In the intraperitoneal infection sepsis model, mice immunized intraperitoneally or subcutaneously with H70+YLN¹⁴ provided significant protection against challenge with three different strains (serotypes 1, 2, and 6A).	234
PspA and PsaA		Intranasal immunization with a combination of PspA and PsaA offered the best protection against nasopharyngeal carriage.	235
PsaA and PspA		Subcutaneous immunization of mice with a vaccine containing the fusion protein PsaA-PspA23 and a single protein, PspA4, significantly increased anti-PspA IgG levels. Mice challenged intranasally with <i>S. pneumoniae</i> from PspA clades 1 to 5 were protected as demonstrated by a decrease in the bacterial load in the lungs and blood along with an increase in survival rates.	236

(Continued)

Table 6. (Continued).

Protein	Results	References
PspA and PdT	Subcutaneous immunization with rPspA-FL-PdT¹⁵ and rPspA-RL-PdT¹⁶ protected mice against intranasal challenge with <i>S. pneumoniae</i> strain A66.1.	237
CbpA and Pneumolysoid	Passive and active immunization with CbpA peptide–L460D fusion protein¹⁷ protected mice from pneumococcal carriage, otitis media, pneumonia, bacteremia, meningitis, and meningococcal sepsis.	238
PdB, PspA, and PspC	Intraperitoneal immunization of mice with a combination of PdB, PspA, and PspC offered the best protection after they were challenged intraperitoneally with virulent type 2 and 6A.	239
PcpA, PhtD, and PlyD1	Immunization with trivalent PPrV containing PcpA, PhtD, and PlyD1 conferred protection in infant mice from lethal pneumonia challenge using serotypes 6A and 3.	240
	Intramuscular vaccination with trivalent PPrV containing PcpA, PhtD, and PlyD1 enhanced the serum and lung antibody levels to the three components, reduced <i>S. pneumoniae</i> lethality, enhanced the clearance of the pathogen via phagocytosis, and reduced lung inflammation and tissue damage.	241
	Intramuscular immunization with trivalent PPrV containing PcpA, PhtD, and PlyD1 protected mice against AOM caused by <i>S. pneumoniae</i>. It reduced the bacterial burden and cytokine inflammatory response in middle ears during AOM induced by <i>S. pneumoniae</i>.	242
	Immunization with PPrV comprising PhtD, PcpA, and PlyD1 induced anti-PhtD and anti-PcpA antibodies in mice that protected against <i>S. pneumoniae</i> through complement- and macrophage-dependent opsonophagocytosis.	243
PsaA and PspA	Subcutaneous immunization of mice with PsaA–PspA fusion protein reduced <i>S. pneumoniae</i> levels in the blood and lungs after intranasal infection. It also protected mice against a fatal challenge with pneumococcal strains expressing different PspAs irrespective of the challenge route.	244
PcsB, StkP, PsaA, and PspA	Subcutaneous immunization of mice with a quadrivalent vaccine formulation containing PcsB, StkP, PsaA, and PspA with adjuvant significantly reduced bacteremia and lung infection when challenged intranasally with <i>S. pneumoniae</i> serotype 1.	245
PiuA and PiaA	Two booster vaccinations consisting of recombinant PiuA and PiaA administered intraperitoneally induced stronger antibody responses (majorly IgG1) in mice than a single or no booster vaccinations.	246
	Intraperitoneal immunization of mice with purified recombinant PiuA and PiaA proteins elicited antibody responses. It also provided a high degree of protection against intraperitoneal challenge with <i>S. pneumoniae</i>.	247
	Intranasal vaccination of mice with PiuA and PiaA generated specific antibody responses in serum and respiratory secretions and protected against intranasal challenge with <i>S. pneumoniae</i>.	248

¹CaP-ΔA146PLY: Calcium phosphate binding domains fused with attenuated pneumolysin mutant, viz., ΔA146PLY.

²ΔA146PLY-SP0148: Fusion protein composed of noncytotoxic mutant of Ply, viz., ΔA146PLY and conserved lipoprotein with high immunogenicity produced by *S. pneumoniae*, viz., SP0148.

³ΔA146 Ply: Ply mutant with one single-amino acid deletion.

⁴PLY-D: Genetic toxoid constructed with two amino acid changes to attenuate PLY toxicity.

⁵PspA-BLP vaccine: PspA proteins from two different families, PspA2 from family 1 and PspA4 from family 2, and formulated with a BLP adjuvant also acting as a carrier.

⁶NP/NCMP PspA4Pro: Poly(glycerol adipate-co-ω-pentadecalactone) polymeric nanoparticles adsorbed with PspA from clade 4 within L-leucine microcarriers.

⁷PspA-nanogel: cCHP nanogel containing PspA.

⁸PspA3+2: Recombinant PspA protein comprising PspA families 1 and 2.

⁹wP: *B. pertussis* vaccine.

¹⁰wP_{low}: Vaccine containing low levels of *B. pertussis* LPS.

¹¹Chitosan-PsaA: Chitosan-DNA nanoparticles expressing PsaA.

¹²PLD: Genetic derivative of Ply.

¹³PlyD6: Nontoxic Ply.

¹⁴H70+YLN: L460D [Ply toxoid with the L460D substitution] fused with protective peptide epitopes from PspC + PspA proline-rich region and SM1 region.

¹⁵rPspA-FL-PdT: Proteins joined by flexible linker.

¹⁶rPspA-RL-PdT: Proteins joined by rigid helix-forming linker.

¹⁷CbpA peptide–L460D fusion protein: Biologically active peptides derived from CbpA of pneumococcus genetically fused to L460D pneumolysoid.

Abbreviations: Ply: Pneumolysin; PlyD1/dPly/PdT: Detoxified pneumolysin; *S. pneumoniae*: *Streptococcus pneumoniae*; IgG: Immunoglobulin G; CBPs: Choline-binding proteins; CbpG: Choline-binding protein G; CbpM: Choline-binding protein M; LytA: N-acetylmuramoyl-L-alanine amidase; LytB: N-acetylglucosaminidase; PspC/CbpA: Pneumococcal surface protein C/Choline-binding protein A; *L. casei*: *Lactobacillus casei*; PspA: Pneumococcal surface protein A; PcpA: Pneumococcal choline binding protein A; CFU: Colony-forming unit; IgA: Immunoglobulin A; BLPs: Bacterium-like particles; SlgA: Secretory immunoglobulin A; RASV: Recombinant attenuated *Salmonella* vaccine; cCHP: Cationic cholesteryl pullulan; *B. pertussis*: *Bordetella pertussis*; LPS: Lipopolysaccharide; PsaA: Pneumococcal surface antigen A; BlyS: B lymphocyte stimulator; IgG1: Immunoglobulin G1; IgG2a: Immunoglobulin G2a; IgG2b: Immunoglobulin G2b; IgG3: Immunoglobulin G3; AOM: Acute otitis media; PHTs: Pneumococcal histidine triad proteins; PhtA: Pneumococcal histidine triad protein A; PhtB: Pneumococcal histidine triad protein B; PhtD: Pneumococcal histidine triad protein D; OMVs: Outer membrane vesicles; PhtE: Pneumococcal histidine triad protein E; PsrP: Pneumococcal serine-rich repeat protein; SRR1: Short serine-rich repeat region; BR: Basic region; Ef-Tu: Elongation factor Tu; PLD: Pneumolysoid; HiB: *Haemophilus influenzae* type b; PiuA: Pneumococcal iron uptake A; Hsp 60: Heat shock protein 60; Hsp 70: Heat shock protein 70; PdB: Pneumolysin toxoid; PPrV: Pneumococcal protein vaccine; PcsB: Putative murine hydrolase; StkP: Serine/threonine kinase protein; PiaA: Pneumococcal iron acquisition A.

determinants.¹⁴⁰ This poses challenges to mounting a protective antibody-mediated immune response as well as explains the difficulty in developing protein-based vaccines.¹⁴⁰ Also, they may not be present in all clinical isolates and may be poorly conserved due to significant sequence heterogeneity.¹⁴⁰ Another drawback of protein antigens is their accessibility to opsonic antibodies.¹⁴⁰ WCVs present a wide range of antigens in their native form and can confer broader protection against *S. pneumoniae*.^{29,102} These vaccines can serve as an economical alternative to PCVs in developing countries because of their low manufacturing costs.^{29,102} WCVs can be developed from killed or

live-attenuated whole-cell or by using genetically modified strains of pneumococcus for ensuring safety and retaining immunogenicity.⁹ Inactivated WCVs offer the advantage of carrying several types of antigens, which are common to all serotypes.¹⁰ These vaccines are relatively safe as the organism is completely inactivated.¹⁰ Live-attenuated vaccines enhance humoral and mucosal immune responses after a single-dose, as compared to inactivated ones.¹⁶⁸ Other merits and demerits of these vaccine technologies are described in detail by Kumraj et al.¹⁶⁹ Table 5 highlights the properties of *S. pneumoniae* proteins that are investigated as vaccine candidates. Tables 6 and 7

Table 7. Results of WCVs investigated in animal models.

Sr. No.	Vaccine candidate	Results	References
Inactivated WCVs			
1.	<i>S. pneumoniae</i> strain RX1 mutated by deleting <i>LytA</i> and killed using ethanol	- Intranasal administration using cholera toxin as an adjuvant conferred protection against nasopharyngeal colonization by serotype 6B in mice as well as conferred protection against illness and death in rats with serotype 3 strains. - In murine models, intranasal administration significantly decreased nasopharyngeal and middle ear colonization by serotypes 6B, 14, and 23F.	249
2.	RM200 (Rx1E PdT Δ <i>lytA</i>): Constructed by replacing the entire <i>LytA</i> gene with a kanamycin resistance gene in strain RX1E and inactivated using beta-propiolactone	Intramuscular immunization in mice offered strong protection against nasopharyngeal colonization with serotype 6B and activated IL-17A priming.	250,251
3.	SPY1: Developed from strain D39 and killed using ethanol	Intranasal immunization with killed SPY1 using cholera toxin as an adjuvant elicited effective protection against colonization with pneumococcal strains 19F and 4 as well as lethal infection with pneumococcal serotypes 2, 3, 14, and 6B.	252
Live-attenuated WCVs			
1.	<i>S. pneumoniae</i> D39 attenuated by removing <i>pep27</i> gene	Intranasal immunization in mice exhibited high levels of IgG and serotype-independent protection against lethal intranasal challenge.	253
2.	Δ <i>pep27</i> Δ <i>comD</i> : Constructed by deleting <i>comD</i> gene	Immunization in mice significantly increased survival time after heterologous challenge and diminished colonization levels independent of serotype.	254
3.	TIGR4 Δ <i>lgt</i> : Constructed by removing <i>Lgt</i> gene in the encapsulated pneumococcal strain TIGR4	- Colonized mouse nasopharynx long enough to induce strong mucosal IgA and IgG2b-dominant systemic antibody responses, exhibiting cross-reactivity to heterologous pneumococcal serotypes. - Intranasal immunization in mice demonstrated serotype-independent protection against pneumococcal challenge.	168
4.	<i>cpsE</i> - <i>endA</i> double mutant strain: Constructed by knocking out <i>cpsE</i> and <i>endA</i> genes in <i>S. pneumoniae</i> D39	Intranasal immunization in mice showed the highest survival rate and longest survival time after a lethal dose intranasal challenge with wild-type <i>S. pneumoniae</i> D39 infection compared to the single-gene knockout strains.	255
5.	Δ <i>cps/piaA</i> and Δ <i>cps/proABC</i> : Strains containing <i>cps</i> locus deletions along with <i>S. pneumoniae</i> virulence factors <i>PsaA</i> or <i>proABC</i>	Previous colonization with these strains in mice protected them against septicemic pneumonia but could not prevent recolonization with the wild-type strain.	256

Abbreviations: WCVs: Whole-cell vaccines; *LytA*: N-acetylmuramoyl-L-alanine amidase; IL-17A: Interleukin-17A; IgG: Immunoglobulin G; *Lgt*: Prolipoprotein diacylglyceryl transferase; IgA: Immunoglobulin A; IgG2b: Immunoglobulin G2b; *PsaA*: Pneumococcal surface antigen A.

Table 8. Results of protein-based vaccines and WCVs (inactivated and live-attenuated) evaluated in clinical trials.

Proteins	Phase	Identifier	Status	Results	References
Protein-based candidates in clinical trials					
PlyD1 ¹	1	NCT01444352	Completed	- Vaccine was immunogenic and safe. - Anti-PlyD1 antibodies produced were shown to be functional in an <i>in vitro</i> assay by neutralizing wild-type Ply activity.	257
PnuBioVax (PlyD6 ² + PspA + PsaA + PiuA + RrgB + RrgA + Hsp60 + Hsp70)	1	NCT02572635	Completed	The vaccine was safe and immunogenic.	233,258
dPly-PhtD $\pm$ PHiD-CV ³	2	NCT00985751	Completed	The vaccine formulations were safe and immunogenic.	259
PhtD	1	NCT01444001	Completed	The vaccine was safe and immunogenic at various doses evaluated.	260
PhtD, PhtD + PcpA	1	NCT01444339	Completed	The vaccine candidates were immunogenic and had promising safety profiles.	261
PcpA + PhtD + PlyD1	1	U1111–1117–7316	Completed	The vaccine was safe and immunogenic.	262
PcpA + PhtD + PlyD1	1	NCT01446926	Completed	NA	263
PhtD	1	NCT01767402	Completed	- The formulations had an acceptable reactogenicity profile, without safety concerns. - Vaccination with PhtD elicited both CD4 T cell and memory B cell responses in adults.	264
dPly, PhtD-dPly $\pm$ PHiD-CV	1	NCT00707798 ⁴	Completed	Vaccine formulations were well-tolerated and immunogenic when given as standalone proteins or combined with PHiD-CV conjugates.	265
PhtD-dPly $\pm$ PCV 8	1	NCT00756067	Completed	Vaccine formulations containing free or conjugated PhtD and dPly had acceptable reactogenicity and safety profiles.	266
PhtD-dPly + PHiD	2	NA	Completed	The vaccine was well-tolerated and immunogenic.	267

(Continued)

Table 8. (Continued).

Proteins	Phase	Identifier	Status	Results	References
PhtD-dPly + PHiD	2	NCT01262872	Completed	• PhtD-dPly + PHiD had no impact on pneumococcal nasopharyngeal carriage. • dPly and PhtD proteins did not alter the immune response to PHiD-CV antigens. • PhtD-dPly + PHiD did not alter the immunogenicity of co-administered routine vaccines.	268,269
PhtD- dPly + PCV13 ⁵	2b	NCT01545375	Completed	• The dPly-PhtD vaccine was immunogenic and well-tolerated.	270
dPLY+PsaA +24CPS ⁶	1	NCT04525599	Completed	• The vaccine had an acceptable safety and tolerability profile. • Immune responses were elicited (IgG concentrations and OPA titers) against 13 common and 10 of the 11 unique <i>S. pneumoniae</i> serotypes.	271
dPLY+PsaA +24CPS	1/2	NCT03803202	Completed	• The vaccine was immunogenic, well-tolerated, and had a safety profile comparable to PCV13.	272
PspA	1	NCT01033409	Completed	NA	273,274
PsaA + StkP + PcsB	1	NCT00873431	Completed	• The vaccine was safe and immunogenic. • It induced protective antibodies against all three proteins. • The vaccine was safe and immunogenic.	275,276
SP_2108 + SP_0148 + SP_1912 (GEN-004)	1	NCT01995617	Completed	• Rise in antigen- specific IgG response was observed.	263,277
SP_2108 + SP_0148 + SP_1912 (GEN-004)	2	NCT02116998	Completed	• Reduced colonization rate by 22–25%.	263,277
WCVs in clinical trials					
Inactivated					
SPWCV developed from strain RM200 (RX1E PdT ΔlytA) using alum as an adjuvant	1	NCT01537185	Completed	• The vaccine was safe, well-tolerated, and elicited a significant rise in IgG responses to PspA and Ply. • An increase in T cell cytokine responses, including IL-17A, was also observed.	278
SPWCV developed from strain RM200 (RX1E PdT ΔlytA) using alum as an adjuvant	1/2	NCT02097472	Completed	NA	29,279,280
SPWCV developed from strain RM200 (RX1E PdT ΔlytA) using alum as an adjuvant	1/2	NCT02543892	Completed	NA	281

¹PlyD1: Detoxified form of Ply.²PlyD6: Non-toxic Ply.³PHiD-CV: 10-valent pneumococcal non-typeable *H. influenzae* protein D conjugate vaccine, viz., Synflorix (GSK).⁴Two groups primed with a formulation containing dPly and PhtD (10 or 30 µg each) continued to the follow-up phase II study (NCT00896064), in which they received a booster dose at 5–9 months after primary vaccination. The study has been completed.²⁸²⁵PCV13 refers to Pfizer's Prevnar 13.⁶dPLY+PsaA +24CPS refers to AFX3772.

Abbreviations: Ply: Pneumolysin; PspA: Pneumococcal surface protein A; PsaA: Pneumococcal surface antigen A; PiuA: Pneumococcal iron uptake A; Hsp 60: Heat shock protein 60; Hsp 70: Heat shock protein 70; dPly: Pneumolysin toxoid; PhtD: Pneumococcal histidine triad protein D; PcpA: Pneumococcal choline binding protein A; CPS: Capsular polysaccharides; IgG: Immunoglobulin G; OPA: Opsonophagocytic activity; *S. pneumoniae*: *Streptococcus pneumoniae*; StkP: Serine/threonine kinase protein; PcsB: Putative murine hydrolase; WCVs: Whole-cell vaccines; SPWCV: *S. pneumoniae* whole-cell vaccine; IL-17A: Interleukin-17A; *H. influenzae*: *Haemophilus influenzae*; GSK: GlaxoSmithKline.

highlight the results of protein-based vaccines and WCVs (inactivated and live-attenuated) investigated in suitable animal models, respectively. Table 8 highlights the results of protein-based vaccines and WCVs (inactivated and live-attenuated) evaluated in clinical trials.

From Tables 6–8, it is evident that there are numerous *S. pneumoniae* proteins, such as pneumolysin (Ply), pneumococcal surface protein A (PspA), pneumococcal surface protein C (PspC), pneumococcal histidine triad protein D (PhtD), pneumococcal surface antigen A (PsaA), etc. that are investigated as vaccine candidates either alone or in combination with other proteins using suitable adjuvants and delivery systems. Some of these proteins have also been explored as carrier proteins to conjugate vaccines. Many of these candidates have shown promising results in animal studies, while some have demonstrated their safety and efficacy in clinical trials. Similarly, many WCVs (inactivated and live-attenuated) have shown promising results in animal studies and clinical trials. However, further studies are

warranted to demonstrate the efficacy and safety of these protein vaccines and WCVs.

Some of the vaccine technologies, such as mRNA vaccines, deoxyribonucleic acid (DNA) vaccines, non-replicating viral vaccines, etc., grabbed the limelight during the coronavirus disease 2019 (COVID-19) pandemic. There exists a possibility that the next generation of pneumococcal vaccines may be developed using such technologies.¹⁰²

Recommendations for future vaccine design and development for LMICs of South Asia

- (1) LMICs should have a robust surveillance system to monitor the pneumococcal serotype prevalence and distribution, the emergence of NVTs, the burden of IPD, and the impact of PCVs. This will give a better and in-depth understanding of the evolving epidemiology of *S. pneumoniae*, aid in formulating and

Table 9. Plausible serotypes that can be incorporated in the existing PCV formulations.

Region	*Plausible serotypes
South Asia	2, 8, 13, 20, 29, 31, 34, 38, 39, 45, 10A, 10F, 11A, 12F, 15A, 15B, 15C, 16F, 17F, 18B, 19B, 22F, 23B, 28A, 28F, 33F, 35B, 35F, 6C, 6D, 7B, 9L, 9N

1) *These serotypes have been implicated in IPD, non-IPD, and drug resistance as shown in Table 2.

Abbreviations: IPD: Invasive pneumococcal disease; Non-IPD: Non-invasive pneumococcal disease.

implementing vaccination policies, and guide the development of next-generation pneumococcal vaccines.

- (2) These countries should have access to high-end and quality diagnostics to understand the AMR pattern of *S. pneumoniae*. Antibiotic stewardship programs and national AMR plans should be religiously implemented. Data from the AMR pattern can play a decisive role in selecting the appropriate serotypes for PCV development.
- (3) Pharmaceutical/Biotech companies should customize or tailor PCVs as per the region-specific prevalence and disease-causing potential of pneumococcal serotypes based on robust surveillance data as well as taking into consideration the pros and cons of adding additional serotypes. For example, companies can develop or modify the existing PCV formulations and incorporate serotypes as shown in Table 9.
- (4) Serotype replacement due to PCVs and regional differences in the dominant serotypes warrant the inclusion of additional serotypes in the existing vaccines. Adding a new serotype will increase the complexity and the cost. This in turn can affect affordability and accessibility in countries with high pneumococcal disease burden and lack of robust healthcare infrastructure. To mitigate this risk, a non-serotype-based approach, such as a protein-based vaccine, a live-attenuated vaccine, or an inactivated vaccine, would be an ideal one. Such vaccines target the highly conserved proteins across all *S. pneumoniae* serotypes and can provide broader protection as compared to the existing PCVs. However, large, robust studies in different geographies are warranted to demonstrate its efficacy and safety in clinical trials as well as the real-world effectiveness of these vaccine candidates.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This research received no external funding.

Notes on contributor

Syed Ahmed is the director and chief executive officer of Techinvention Lifecare Private Limited, an innovation-driven biotech company focused

on vaccines, diagnostics, and biotherapeutics for infectious diseases. He is also the board member and vice president of Emerging Biopharmaceutical Manufacturers Network (EBPMN), a not-for-profit pan global association supporting access and manufacturing of biopharmaceuticals across low- and middle-income countries (LMICs). Syed is also the co-founder of another not-for-profit organization in its inception stage, namely Association for Veterinary Vaccine Manufacturers (AVVM), with a focus on enabling access and equity of veterinary vaccines in LMICs. Syed is the co-inventor and has 15+ patents filed in the field of discovery and development of immunobiologicals. He's also the corresponding author of four manuscripts in the arena of vaccines and biotherapeutics published in global journals of repute. He's also leading quite a few 'One Health' initiatives and focused on interventions in the domain of infectious diseases that can impact climate change and sustainability.

Author contributions

Conceptualization, P.D., S.S., K.S., and S.A.; writing – original draft preparation, P.D., S.S., K.S., V.K., A.B., S.D., P.S. S.A.; writing – review and editing, P.D., S.S., D.S., and S.A. All authors have read and agreed to the published version of the manuscript.

References

- Leonard A, Möhlis K, Schlüter R, Taylor E, Lalk M, Methling K. Exploring metabolic adaptation of *Streptococcus pneumoniae* to antibiotics. *J Antibiot (Tokyo)*. 2020;73(7):441–454. doi:10.1038/s41429-020-0296-3.
- Plainvert C, Varon E, Viriot D, Kempf M, Plainvert C, Alauzet C, Auger G, Batah J, Brieu N, Cattoir V, et al. Invasive pneumococcal infections in France: changes from 2009 to 2021 in antibiotic resistance and serotype distribution of *Streptococcus pneumoniae* based on data from the French Regional Pneumococcal Observatories network. *Infect Dis Now*. 2023;53(1):104632. doi:10.1016/j.idnow.2022.11.001.
- Deloria Knoll M, Bennett J, Garcia Quesada M, Kagucia E, Peterson M, Feikin D, Cohen A, Hetrich M, Yang Y, Sinkevitch J, et al. Global landscape review of serotype-specific invasive pneumococcal disease surveillance among countries using PCV10/13: the pneumococcal serotype replacement and distribution estimation (PSERENADE) project. *Microorganisms*. 2021;9(4):742. doi:10.3390/microorganisms9040742.
- Cui YA, Patel H, Wm O, Li S, Saddier P. Pneumococcal serotype distribution: a snapshot of recent data in pediatric and adult populations around the world. *Hum Vaccines Immunother*. 2017;13(6):1229–1241. doi:10.1080/21645515.2016.1277300.
- Teixeira R, Kossyvakı V, Galvez P, Méndez C. Pneumococcal serotype evolution and burden in European adults in the last decade: a systematic review. *Microorganisms*. 2023;11(6):1376. doi:10.3390/microorganisms11061376.
- Kim SH, Chung DR, Song J-H, Baek JY, Thamlikitkul V, Wang H, Carlos C, Ahmad N, Arushothy R, Tan SH, et al. Changes in serotype distribution and antimicrobial resistance of *Streptococcus pneumoniae* isolates from adult patients in Asia: emergence of drug-resistant non-vaccine serotypes. *Vaccine*. 2020;38(38):6065–6073. doi:10.1016/j.vaccine.2019.09.065.
- Pneumococcal Disease Surveillance and Trends. [accessed 2024 Jul 3]. <https://www.cdc.gov/pneumococcal/php/surveillance/index.html>.
- Cillóniz C, García-Vidal C, Ceccato A, Torres A. Antimicrobial resistance among *Streptococcus pneumoniae*. In: Fong I; Shlaes D Drlica K, editors. *Antimicrobial Resistance in the 21st Century*. Cham: Springer International Publishing; 2018. p. 13–38.
- Li S, Liang H, Zhao S-H, Yang X-Y, Guo Z. Recent progress in pneumococcal protein vaccines. *Front Immunol*. 2023;14:1278346. doi:10.3389/fimmu.2023.1278346.
- Silva PH, Vázquez Y, Campusano C, Retamal-Díaz A, Lay MK, Muñoz CA, González PA, Kalergis AM, Bueno SM. Non-capsular based immunization approaches to prevent *Streptococcus*

- pneumoniae infection. *Front Cell Infect Microbiol.* 2022;12:949469. doi:10.3389/fcimb.2022.949469.
11. Weiser JN, Ferreira DM, Paton JC. Streptococcus pneumoniae: transmission, colonization and invasion. *Nat Rev Microbiol.* 2018;16(6):355–367. doi:10.1038/s41579-018-0001-8.
 12. Micoli F, Romano MR, Carboni F, Adamo R, Berti F. Strengths and weaknesses of pneumococcal conjugate vaccines. *Glycoconj J.* 2023;40(2):135–148. doi:10.1007/s10719-023-10100-3.
 13. Rodgers GL, Whitney CG, Klugman KP. Triumph of pneumococcal conjugate vaccines: overcoming a common foe. *J Infect Dis.* 2021;224(Supplement_4):S352–9. doi:10.1093/infdis/jiaa535.
 14. Kartasmita CB, Rezeki Hadinegoro S, Kurniati N, Triasih R, Halim C, Gamil G. Triumph of pneumococcal conjugate vaccines: overcoming a common foe. *Infect Dis Ther.* 2020;9(4):723–736. doi:10.1007/s40121-020-00330-5.
 15. Løchen A, Croucher NJ, Anderson RM. Divergent serotype replacement trends and increasing diversity in pneumococcal disease in high income settings reduce the benefit of expanding vaccine valency. *Sci Rep.* 2020;10(1):18977. doi:10.1038/s41598-020-75691-5.
 16. Song JY, Nahm MH, Moseley MA. Clinical implications of pneumococcal serotypes: invasive disease potential, clinical presentations, and antibiotic resistance. *J Korean Med Sci.* 2013;28(1):4. doi:10.3346/jkms.2013.28.1.4.
 17. Wattal C, Goel N, Byotra SP. pneumococcal serotypes in adults ≥50 years of age. *Indian J Med Microbiol.* 2017;35(1):95–100. doi:10.4103/ijmm.IJMM_16_132.
 18. Du Q, Shi W, Yu D, Yao K. Epidemiology of non-vaccine serotypes of Streptococcus pneumoniae before and after universal administration of pneumococcal conjugate vaccines. *Hum Vaccines Immunother.* 2024;17(12):5628–5637. doi:10.1080/21645515.2021.1985353.
 19. Grant LR, Slack MPE, Theilacker C, Vojcic J, Dion S, Reinert R-R, Jodar L, Gessner BD. Distribution of serotypes causing invasive pneumococcal disease in children from high-income countries and the impact of pediatric pneumococcal vaccination. *Clin Infect Dis.* 2023;76(3):e1062–70. doi:10.1093/cid/ciac475.
 20. Lister AJJ, Le CF, Cheah ESG, Desa MNM, Cleary DW, Clarke SC. Serotype distribution of invasive, non-invasive and carried Streptococcus pneumoniae in Malaysia: a meta-analysis. *Pneumonia.* 2021;13(1):9. doi:10.1186/s41479-021-00086-7.
 21. El-Beleidy A, El-Saied M, Fasseeh N, El Saie RZ, Haridy H. A Systematic review of pneumococcal carriage, disease, antimicrobial resistance, and vaccination in Egyptian children aged 18 years and younger. *Infect Dis Ther.* 2021;10(4):2119–2155. doi:10.1007/s40121-021-00523-6.
 22. Abebaw T-A, Aregay WK, Ashami MT. Risk factors for childhood pneumonia at Adama Hospital Medical College, Adama, Ethiopia: a case-control study. *Pneumonia Nathan Qld.* 2022;14(1):9. doi:10.1186/s41479-022-00102-4.
 23. Jaiswal N, Singh M, Thumburu KK, Bharti B, Agarwal A, Kumar A, Kaur H, Chadha N. Burden of invasive pneumococcal disease in children aged 1 month to 12 years living in South Asia: A Systematic Review. *PLOS ONE.* 2014;9(5):e96282. doi:10.1371/journal.pone.0096282.
 24. Wasserman MD, Perdrizet J, Grant L, Hayford K, Singh S, Saharia P, Horn EK, Farkouh RA. Clinical and economic burden of pneumococcal disease due to serotypes contained in current and investigational pneumococcal conjugate vaccines in children under five years of age. *Infect Dis Ther.* 2021;10(4):2701–2720. doi:10.1007/s40121-021-00544-1.
 25. Chan J, Lai JY, Von Mollendorf C, Blyth C, Dance DAB, Datta S, Dunne EM, Ford R, Fox K, Hinds J, et al. Determining the pneumococcal conjugate vaccine coverage required for indirect protection within Asia and the Pacific: a prospective observational study. *Lancet Glob Health.* 2019;7:S15. doi:10.1016/S2214-109X(19)30100-7.
 26. View-hub Report: Global Vaccine Introduction and Implementation. [accessed 2025 Jan 18]. https://view-hub.org/sites/default/files/2024-09/VIEW-hub_Report_August2024.pdf.
 27. Syeed MS, Ghule P, Le LM, Veettil SK, Horn EK, Perdrizet J, Wasserman M, Thakkinstian A, Chaiyakunapruk N. Pneumococcal vaccination in children: a systematic review and meta-analysis of cost-effectiveness studies. *Value Health.* 2023;26(4):598–611. doi:10.1016/j.jval.2022.10.006.
 28. Chan J, Nguyen CD, Dunne EM, Kim Mulholland E, Mungun T, Pomat WS, Rafai E, Satzke C, Weinberger DM, Russell FM. Using pneumococcal carriage studies to monitor vaccine impact in low- and middle-income countries. *Vaccine.* 2019;37(43):6299–6309. doi:10.1016/j.vaccine.2019.08.073.
 29. Masomian M, Ahmad Z, Ti Gew L, Poh CL. Development of Next Generation Streptococcus pneumoniae Vaccines Conferring Broad Protection. *Vaccines.* 2020;8(1):132. doi:10.3390/vaccines8010132.
 30. Van Tonder AJ, Gladstone RA, Lo SW, Nahm MH, du Plessis M, Cornick J, Kwambana-Adams B, Madhi SA, Hawkins PA, Benisty R, et al. Putative novel cps loci in a large global collection of pneumococci. *Microb Genomics.* 2019;5(7). doi:10.1099/mgen.0.000274.
 31. Manzanal A, Vicente D, Alonso M, Azkue N, Ercibengoa M, Marimón JM. Impact of the progressive uptake of pneumococcal conjugate vaccines on the epidemiology and antimicrobial resistance of invasive pneumococcal disease in Gipuzkoa, northern Spain, 1998–2022. *Front Public Health.* 2023;11:1238502. doi:10.3389/fpubh.2023.1238502.
 32. Weinberger DM, Warren JL, Dalby T, Shapiro ED, Valentiner-Branth P, Slotved H-C, Harboe ZB. Differences in the impact of pneumococcal serotype replacement in individuals with and without underlying medical conditions. *Clin Infect Dis Off Publ Infect Dis Soc Am.* 2019;69(1):100–106. doi:10.1093/cid/ciy875.
 33. Lewnard JA, Hanage WP. Making sense of differences in pneumococcal serotype replacement. *Lancet Infect Dis.* 2019;19(6):e213–20. doi:10.1016/S1473-3099(18)30660-1.
 34. Wambugu P, Shah M-M, Nguyen H-A, Le K-A, Le H-H, Vo H-M, Toizumi M, Bui M-X, Dang D-A, Yoshida L-M. Molecular epidemiology of streptococcus pneumoniae detected in hospitalized pediatric acute respiratory infection cases in Central Vietnam. *Pathogens.* 2023;12(7):943. doi:10.3390/pathogens12070943.
 35. Sabbar H, Mahraoui C, Bastias García M, Jroundi I. Streptococcus pneumoniae vaccination strategies and its expected impact on penicillin non-susceptibility in children under the age of five: let's recap! *Vaccine X.* 2022;11:100170. doi:10.1016/j.jvacx.2022.100170.
 36. Lo SW, Gladstone RA, Van Tonder AJ, Lees JA, du Plessis M, Benisty R, Givon-Lavi N, Hawkins PA, Cornick JE, Kwambana-Adams B, et al. Pneumococcal lineages associated with serotype replacement and antibiotic resistance in childhood invasive pneumococcal disease in the post-PCV13 era: an international whole-genome sequencing study. *Lancet Infect Dis.* 2019;19(7):759–769. doi:10.1016/S1473-3099(19)30297-X.
 37. Balsells E, Guillot L, Nair H, Kyaw MH, Borrow R. Serotype distribution of Streptococcus pneumoniae causing invasive disease in children in the post-pcv era: a systematic review and meta-analysis. *PLOS ONE.* 2017;12(5):e0177113. doi:10.1371/journal.pone.0177113.
 38. Scott NR, Mann B, Tuomanen EI, Orihuela CJ. Multi-valent protein hybrid pneumococcal vaccines: a strategy for the next. *Vaccines.* 2021;9(3):209. doi:10.3390/vaccines9030209.
 39. Zabihullah R, Dhoubhadel BG, Rauf FA, Shafiq SA, Suzuki M, Watanabe K, Yoshida LM, Yasunami M, Zabihullah S, Parry CM, et al. Risk for death among children with pneumonia, Afghanistan. *Emerg Infect Dis.* 2017;23(8):1404–1408. doi:10.3201/eid2308.151550.
 40. Pichichero M, Malley R, Kaur R, Zagursky R, Anderson P. Acute otitis media pneumococcal disease burden and nasopharyngeal colonization in children due to serotypes included and not included in current and new pneumococcal conjugate vaccines. *Expert Rev Vaccines.* 2023;22(1):118–138. doi:10.1080/14760584.2023.2162506.

41. Malaker R, Saha S, Hanif M, Ahmed A, Saha S, Hasanuzzaman M, Khondakar T, Islam M, Baqui AH, Santosham M, et al. Invasive pneumococcal infections in children with nephrotic syndrome in Bangladesh. *Pediatr Infect Dis J*. 2019;38(8):798–803. doi:10.1097/INF.0000000000002386.
42. Baqui AH, McCollum ED, Mahmud A, Roy A, Chowdhury NH, Rafiqullah I, Rizvi SJR, Begum N, Mitra DK, Khanam R, et al. Population-based incidence and serotype distribution of invasive pneumococcal disease prior to introduction of conjugate pneumococcal vaccine in Bangladesh. *PLOS ONE*. 2020;15(2):e0228799. doi:10.1371/journal.pone.0228799.
43. Saha SK, Hossain B, Islam M, Hasanuzzaman M, Saha S, Hasan M, Darmstadt GL, Chowdhury M, Arifeen SE, Baqui AH, et al. Epidemiology of invasive pneumococcal disease in Bangladeshi children before introduction of pneumococcal conjugate vaccine. *Pediatr Infect Dis J*. 2016;35(6):655–661. doi:10.1097/INF.0000000000001037.
44. Siddik AB, Tanvir NA, Bhuyan GS, Alam MS, Islam Z, Bulbul MRH, Moniruzzaman M, Halder CE, Rahman T, Endtz H, et al. Bacterial and viral etiology of acute respiratory infection among the Forcibly Displaced Myanmar Nationals (FDMNs) in fragile settings in Cox's Bazar- a prospective case-control study. *PLoS Negl Trop Dis*. 2023;17(4):e0011189. doi:10.1371/journal.pntd.0011189.
45. Naziat H, Saha S, Islam M, Saha S, Uddin MJ, Hussain M, Luby SP, Darmstadt GL, Whitney CG, Gessner BD, et al. Epidemiology of Otitis media with Otorrhea among Bangladeshi children: Baseline study for future assessment of pneumococcal conjugate vaccine impact. *Pediatr Infect Dis J*. 2018;37(7):715–721. doi:10.1097/INF.0000000000002077.
46. Apte A, Dayma G, Naziat H, Williams L, Sanghavi S, Uddin J, Kawade A, Islam M, Kar S, Li Y, et al. Nasopharyngeal pneumococcal carriage in South Asian infants: results of observational cohort studies in vaccinated and unvaccinated populations. *J Glob Health*. 2021;11:04054. doi:10.7189/jogh.11.04054.
47. Vidanapathirana G, Angulmaduwa S, Munasinghe T, Ekanayake A, Kudagammana T, Dissanayaka N, Liyanapathirana V. Pneumococcal colonization among healthy and hospitalized vaccine-naïve Sri Lankan children. *Vaccine*. 2020;38(46):7308–7315. doi:10.1016/j.vaccine.2020.09.040.
48. Vidanapathirana G, Angulmaduwa ALSK, Munasinghe TS, Ekanayake EWMA, Harasgama P, Kudagammana ST, Dissanayake BN, Liyanapathirana LVC. Comparison of pneumococcal colonization density among healthy children and children with respiratory symptoms using real time PCR (RT-PCR). *BMC Microbiol*. 2022;22(1):31. doi:10.1186/s12866-022-02442-z.
49. Carter MJ, Gurung M, Pokhrel B, Bijukchhe SM, Karmacharya S, Khadka B, Maharjan A, Bhattarai S, Shrestha S, Khadka B, et al. Childhood invasive bacterial disease in Kathmandu, Nepal (2005–2013). *Pediatr Infect Dis J*. 2022;41(3):192–198. doi:10.1097/INF.0000000000003421.
50. Shrestha S, Gurung M, Amatya P, Bijukchhe S, Bose AS, Carter MJ, Gautam MC, Gurung S, Hinds J, Kandasamy R, et al. Effect of the of 10-valent pneumococcal conjugate vaccine in Nepal 4 years after introduction: an observational cohort study. *Lancet Glob Health*. 2022;10(10):e1494–504. doi:10.1016/S2214-109X(22)00281-9.
51. Kandasamy R, Gurung M, Thapa A, Ndimah S, Adhikari N, Murdoch DR, Kelly DF, Waldron DE, Gould KA, Thorson S, et al. Multi-serotype pneumococcal nasopharyngeal carriage prevalence in vaccine naïve Nepalese children, assessed using molecular serotyping. *PLOS ONE*. 2015;10(2):e0114286. doi:10.1371/journal.pone.0114286.
52. Kandasamy R, Lo S, Gurung M, Carter MJ, Gladstone R, Lees J, Shrestha S, Thorson S, Bijukchhe S, Gautam MC, et al. Effect of childhood vaccination and antibiotic use on pneumococcal populations and genome-wide associations with disease among children in Nepal: an observational study. *The Lancet Microbe*. 2022;3(7):e503–11. doi:10.1016/S2666-5247(22)00066-0.
53. Jullien S, Sharma R, Lhamu Mynak M, Henares D, Muñoz-Almagro C, Bassat Q. Pneumococcal nasopharyngeal carriage among Bhutanese children hospitalized with clinical pneumonia: serotypes and viral co-infection. *BMC Infect Dis*. 2020;20(1):940. doi:10.1186/s12879-020-05674-4.
54. Shahid S, Nisar MI, Jehan F, Ahmed S, Kabir F, Hotwani A, Muneer S, Qazi MF, Muhammad S, Ali A, et al. Co-carriage of *Staphylococcus aureus* and *Streptococcus pneumoniae* among children younger than 2 years of age in a rural population in Pakistan. *Clin Epidemiol Glob Health*. 2023;21:101293. doi:10.1016/j.cegh.2023.101293.
55. Riaz A, Mohiuddin S, Husain S, Yousafzai MT, Sajid M, Kabir F, Rehman NU, Mirza W, Salam B, Nadeem N, et al. Effectiveness of 10-valent pneumococcal conjugate vaccine against vaccine-type invasive pneumococcal disease in Pakistan. *Int J Infect Dis*. 2019;80:28–33. doi:10.1016/j.ijid.2018.12.007.
56. Nisar MI, Nayani K, Akhund T, Riaz A, Irfan O, Shakoor S, Muneer S, Muslim S, Hotwani A, Kabir F, et al. Nasopharyngeal carriage of *Streptococcus pneumoniae* in children under 5 years of age before introduction of pneumococcal vaccine (PCV10) in urban and rural districts in Pakistan. *BMC Infect Dis*. 2018;18(1):672. doi:10.1186/s12879-018-3608-5.
57. Khan F, Khan MA, Ahmed N, Bashir H, Taahir S, Zafar AU. Molecular Characterization of Pneumococcal Surface Protein A (PspA), Serotype Distribution and Antibiotic Susceptibility of *Streptococcus pneumoniae* Strains Isolated from Pakistan. *Infect Dis Ther*. 2018 Jun;7(2):291. doi:10.1007/s40121-018-0199-9.
58. Nisar MI, Jehan F, Shahid S, Ahmed S, Shakoor S, Kabir F, Hotwani A, Muneer S, Khalid F, Muhammad S, et al. Serotype-specific effectiveness against pneumococcal carriage and serotype replacement after ten-valent Pneumococcal Conjugate Vaccine (PCV10) introduction in Pakistan. *PLOS ONE*. 2022;17(1):e0262466. doi:10.1371/journal.pone.0262466.
59. Shakoor S, Kabir F, Khowaja AR, Qureshi SM, Jehan F, Qamar F, Whitney CG, Zaidi AKM. Pneumococcal serotypes and serogroups causing invasive disease in Pakistan, 2005–2013. *PLOS ONE*. 2014;9(6):e98796. doi:10.1371/journal.pone.0098796.
60. Nisar MI, Shahid S, Jehan F, Ahmed S, Shakoor S, Kabir F, Hotwani A, Munir S, Khalid F, Muhammad S, et al. Antimicrobial resistance in pneumococcal carriage isolates from children under 2 years of age in rural Pakistan. *Microbiol Spectr*. 2021;9(3):e01019–21. doi:10.1128/Spectrum.01019-21.
61. Javaid N, Lo SW, Nisar MI, Basharat A, Jaleel H, Rasool K, Sultana Q, Kabir F, Hotwani A, Breiman RF, et al. Strain features of pneumococcal isolates in the pre- and post-PCV10 era in Pakistan. *Microb Genomics*. 2024;10(1). doi:10.1099/mgen.0.001163.
62. Shahid S, Khan A, Nisar MI, Khalid F, Qazi MF, Ahmed S, Kabir F, Hotwani A, Muneer S, Ali SA, et al. Pneumococcal carriage in infants post-PCV10 introduction in Pakistan: results from serial cross-sectional surveys. *Vaccines*. 2022;10(6):971. doi:10.3390/vaccines10060971.
63. Nisar MI, Ahmed S, Jehan F, Shahid S, Shakoor S, Kabir F, Hotwani A, Munir S, Muhammad S, Khalid F, et al. Direct and indirect effect of 10 valent pneumococcal vaccine on nasopharyngeal carriage in children under 2 years of age in Matiari, Pakistan. *Vaccine*. 2021;39(8):1319–1327. doi:10.1016/j.vaccine.2020.12.066.
64. Jaiswal N, Singh M, Das RR, Jindal I, Agarwal A, Thumburu KK, Kumar A, Chauhan A. Distribution of serotypes, vaccine coverage, and antimicrobial susceptibility pattern of *Streptococcus pneumoniae* in children living in SAARC countries: A systematic review. *PLOS ONE*. 2014;9(9):e108617. doi:10.1371/journal.pone.0108617.
65. Dananché C, Paranhos-Baccalà G, Messaoudi M, Sylla M, Awasthi S, Bavdekar A, Sanghavi S, Diallo S, Pape JW, Rouzier V, et al. Serotypes of *Streptococcus pneumoniae* in Children Aged <5 Years Hospitalized With or Without Pneumonia in Developing and Emerging Countries: A Descriptive, Multicenter Study. *Clin Infect Dis*. 2020;70(5):875–883. doi:10.1093/cid/ciz277.
66. Balaji V, Jayaraman R, Verghese V, Baliga P, Kurien T. Pneumococcal serotypes associated with invasive disease in under five children in India & implications for vaccine policy.

- Indian J Med Res. 2015;142(3):286. doi:10.4103/0971-5916.166588.
67. Veeraraghavan B, Jayaraman R, John J, Varghese R, Neeravi A, Verghese VP, Thomas K. Customized sequential multiplex PCR for accurate and early determination of invasive pneumococcal serotypes found in India. *J Microbiol Methods*. 2016;130:133–135. doi:10.1016/j.mimet.2016.09.007.
 68. Nagaraj S, Kalal BS, Manoharan A, Shet A. Streptococcus pneumoniae serotype prevalence and antibiotic resistance among young children with invasive pneumococcal disease: experience from a tertiary care center in South India. *GERMS*. 2017;7(2):78–85. doi:10.18683/germs.2017.1112.
 69. Singh J, Sundaresan S, Manoharan A, Shet A. Serotype distribution and antimicrobial susceptibility pattern in children $\leq$ 5 years with invasive pneumococcal disease in India – a systematic review. *Vaccine*. 2017;35(35):4501–4509. doi:10.1016/j.vaccine.2017.06.079.
 70. Jayaraman R, Varghese R, Kumar JL, Neeravi A, Shanmugasundaram D, Ralph R, Thomas K, Veeraraghavan B. Invasive pneumococcal disease in Indian adults: 11 years' experience. *J Microbiol Immunol Infect*. 2019;52(5):736–742. doi:10.1016/j.jmii.2018.03.004.
 71. Sutcliffe CG, Shet A, Varghese R, Veeraraghavan B, Manoharan A, Wahl B, Chandy S, Sternal J, Khan R, Singh RK, et al. Nasopharyngeal carriage of Streptococcus pneumoniae serotypes among children in India prior to the introduction of pneumococcal conjugate vaccines: a cross-sectional study. *BMC Infect Dis*. 2019;19(1):605. doi:10.1186/s12879-019-4254-2.
 72. Manoharan A, Manchanda V, Balasubramanian S, Lalwani S, Modak M, Bai S, Vijayan A, Shet A, Nagaraj S, Karande S, et al. Invasive pneumococcal disease in children aged younger than 5 years in India: a surveillance study. *Lancet Infect Dis*. 2017;17(3):305–312. doi:10.1016/S1473-3099(16)30466-2.
 73. Gupta P, Awasthi S, Gupta U, Verma N, Rastogi T, Pandey AK, Naziat H, Rahman H, Islam M, Saha S. Nasopharyngeal carriage of streptococcus pneumoniae serotypes among healthy children in Northern India. *Curr Microbiol*. 2023;80(1):41. doi:10.1007/s00284-022-03114-x.
 74. John J, Varghese R, Lionell J, Neeravi A, Veeraraghavan B. Nonvaccine pneumococcal serotypes among children with invasive pneumococcal disease. *Indian Pediatr*. 2018;55(10):874–876. doi:10.1007/s13312-018-1398-x.
 75. Verghese VP, Veeraraghavan B, Jayaraman R, Varghese R, Neeravi A, Jayaraman Y, Thomas K, Mehendale SM. Increasing incidence of penicillin- and cefotaxime-resistant streptococcus pneumoniae causing meningitis in India: time for revision of treatment guidelines? *Indian J Med Microbiol*. 2017;35(2):228–236. doi:10.4103/ijmm.IJMM_17_124.
 76. Sharma S, Sharma M, Ray P, Chakraborti A. Antimicrobial susceptibility pattern and serotype distribution of streptococcus pneumoniae isolates from a hospital-based study in Chandigarh, North India. *Cureus*. 2022; published online Jan 19. DOI:10.7759/cureus.21437.
 77. Vandana G, Feroze AG, Geetha N, Avid H, Ravi Kumar KL. Pan India distribution of pneumococcal serotypes (PIDOPS) causing invasive pneumococcal disease and pneumonia in children between 6 weeks and 5 years and their antimicrobial resistance – Phase I. *Pediatr Infect Dis*. 2016;8(2):47–51. doi:10.1016/j.pid.2016.06.004.
 78. Balasubramanian S, Singh P, Dhanalakshmi K, Putlibai S, Varghese R. Clinico-microbiological profile of invasive pneumococcal disease amongst hospitalised children in South India. *Indian J Med Microbiol*. 2019;37(4):598–599. doi:10.4103/ijmm.IJMM_20_135.
 79. Jayaraman Y, Veeraraghavan B, Girish Kumar CP, Sukumar B, Rajkumar P, Kangusamy B, Verghese VP, Varghese R, Jayaraman R, Kapoor AN, et al. Hospital-based sentinel surveillance for bacterial meningitis in under-five children prior to the introduction of the PCV13 in India. *Vaccine*. 2021;39(28):3737–3744. doi:10.1016/j.vaccine.2021.05.041.
 80. Verma N, Gupta P, Pandey AK, Awasthi S. Nasopharyngeal carriage of Streptococcus pneumoniae serotypes among sick and healthy children in Northern India: a case-control study. *Vaccine*. 2023;41(44):6619–6624. doi:10.1016/j.vaccine.2023.09.029.
 81. Manoharan A, Jayaraman R. Pneumococcal aaccines. *Indian J Med Microbiol*. 2018;36(4):465–474. doi:10.4103/ijmm.IJMM_18_442.
 82. Mazumdar D, Sarma A, Medhi D, Dutta R, Kataki M, Baishya L, Dutta BS, Saikia L. Capsular typing of Streptococcus pneumoniae isolated from clinical specimens in Gauhati Medical College and hospital, Assam, India. *Indian J Med Microbiol*. 2023;44:100350. doi:10.1016/j.ijmm.2023.01.008.
 83. Varghese R, Neeravi A, Subramanian N, Pavithra B, Kavipriya A, Kumar JL, Kumar CPG, Jeyraman Y, Karthik G, Verghese VP, et al. Clonal similarities and sequence-Type diversity of invasive and carriage Streptococcus pneumoniae in India among children under 5 years. *Indian J Med Microbiol*. 2019;37(3):358–361. doi:10.4103/ijmm.IJMM_19_348.
 84. Nagaraj G, Govindan V, Ganaie F, Venkatesha VT, Hawkins PA, Gladstone RA, McGee L, Breiman RF, Bentley SD, Klugman KP, et al. Streptococcus pneumoniae genomic datasets from an Indian population describing pre-vaccine evolutionary epidemiology using a whole genome sequencing approach. *Microb Genomics*. 2021;7(9). doi:10.1099/mgen.0.000645.
 85. Napoleon M, Rosemol V, John M, Varghese AM, Periyasamy J, Balaji V, Naina P. Nasopharyngeal colonization of otopathogens in South Indian children with acute otitis media – a case control pilot study. *J Otol*. 2021;16(4):220–224. doi:10.1016/j.joto.2021.02.004.
 86. Jain S, Das BK, Mahajan N, Kapil A, Chaudhry R, Sood S, Kabra SK, Dwivedi SN. Molecular capsular typing and multi locus sequence typing of invasive, non-invasive and commensal Streptococcus pneumoniae isolates from North India. *Indian J Med Microbiol*. 2020;38(1):78–86. doi:10.4103/ijmm.IJMM_20_111.
 87. Varghese R, Neeravi A, Jacob JJ, Vasudevan K, Kumar JL, Subramanian N, Veeraraghavan B. Comparative genomics of invasive Streptococcus pneumoniae CC320/271 serotype 19F/19A before the introduction of pneumococcal vaccine in India. *Mol Biol Rep*. 2021;48(4):3265–3276. doi:10.1007/s11033-021-06353-6.
 88. Jayaraman Y, Veeraraghavan B, Chethrapilly Purushothaman GK, Sukumar B, Kangusamy B, Nair Kapoor A, Gupta N, Mehendale SM. Burden of bacterial meningitis in India: Preliminary data from a hospital based sentinel surveillance network. *PLOS ONE*. 2018;13(5):e0197198. doi:10.1371/journal.pone.0197198.
 89. Rafiqullah IM, Varghese R, Hellmann KT, Velmurugan A, Neeravi A, Kumar Daniel JL, Vidal JE, Kompithra RZ, Verghese VP, Veeraraghavan B, et al. Pneumococcal population genomics changes during the early time period of conjugate vaccine uptake in southern India. *Microb Genomics*. 2024;10(2). doi:10.1099/mgen.0.001191.
 90. Kumar S, Purakayastha DR, Kapil A, Saha S, Dawood FS, Das BK, Amarchand R, Kumar R, Lafond KE, Jain S, et al. Carriage rates and antimicrobial sensitivity of pneumococci in the upper respiratory tract of children less than ten years old, in a North Indian rural community. *PLOS ONE*. 2021;16(2):e0246522. doi:10.1371/journal.pone.0246522.
 91. Arya BK, Bhattacharya SD, Sutcliffe CG, Ganaie F, Bhaskar A, Bhattacharyya S, Niyogi SK, Moss WJ, Panda S, Ravikumar KL, et al. Nasopharyngeal pneumococcal colonization and impact of a single dose of 13-valent pneumococcal conjugate vaccine in Indian children with HIV and their unvaccinated parents. *Pediatr Infect Dis J*. 2018;37(5):451–458. doi:10.1097/INF.0000000000001800.
 92. Ganaie F, Nagaraj G, Govindan V, Basha R, Hussain M, Ashraf N, Ahmed S, Ravi Kumar KL. Impact of Hajj on the S. pneumoniae carriage among Indian pilgrims during 2016- a longitudinal molecular surveillance study. *Travel Med Infect Dis*. 2018;23:64–71. doi:10.1016/j.tmaid.2018.04.002.
 93. Samson D, Rupa V, Veeraraghavan B, Varghese R, Isaac R, J L. Follow up of a birth cohort to identify prevalence and risk factors for otitis media among Indian children in the eighth year of life. *Int J Pediatr Otorhinolaryngol*. 2020;137:110201. doi:10.1016/j.ijporl.2020.110201.

94. Arya BK, Bhattacharya SD, Harigovind G, Das RS, Khan T, Ganaie F, Niyogi SK, Ravikumar KL, Manoharan A, Bhattacharyya S, et al. Streptococcus pneumoniae acquisition and carriage in vaccine Naïve Indian children with HIV and their parents: a longitudinal-household study. *Indian J Pediatr.* 2019;86(11):1002–1010. doi:10.1007/s12098-019-02995-4.
95. Veeraraghavan B, Varghese R, Saigal K, Balasubramanian S, Bai PSP, Lal Y B, Neeravi A, Baskar P, Anandhan K, Kumar CPG, et al. Activity of novel lactone ketolide nafithromycin against multi-centric invasive and non-invasive pneumococcal isolates collected in India. *JAC-Antimicrob Resist.* 2021;3(2):dlab066. doi:10.1093/jacamr/dlab066.
96. Kanungo R, Anandhalakshmi S, Sheeladevi C, Sudhagar M, John J, Prashanth K. Fatal infection in adults by pneumolysin & autolysin producing, non-vaccine serotype Streptococcus pneumonia. *Indian J Med Res.* 2016;143(4):514. doi:10.4103/0971-5916.184307.
97. Raman R, Sankar J, Putlibai S, Raghavan V. Demographic profile of healthy children with Nasopharyngeal colonisation of Streptococcus pneumoniae: a research paper. *Indian J Med Microbiol.* 2017;35(4):607–609. doi:10.4103/ijmm.IJMM_15_347.
98. Amritha GN, M N, R APS, Shanmugam P, Jayaraman P. A comparative profile of Oropharyngeal colonization of Streptococcus pneumoniae and Hemophilus influenzae among HealthCare Workers (HCW) in a tertiary care hospital and non-healthcare individuals. *J Prev Med Hyg.* 2020;61(3):E 379. doi:10.15167/2421-4248/jpmh2020.61.3.1479.
99. Arjun R, Ratheesh RS, Mohan V, Uduman S, Jalaludeen S, Prabhakaran A, Sasidharan A, Niyas VKM. Susceptibility and serotypes of Streptococcus pneumoniae isolates in invasive pneumococcal disease: a study from Kerala, South India. *Infez Med.* 2020;28(4):558–564.
100. Varghese R, Daniel JL, Neeravi A, Baskar P, Manoharan A, Sundaram B, Manchanda V, Saigal K, Yesudhasan BL, Veeraraghavan B. Multicentric analysis of Erythromycin resistance determinants in invasive Streptococcus pneumoniae; associated serotypes and sequence types in India. *Curr Microbiol.* 2021;78(8):3239–3245. doi:10.1007/s00284-021-02594-7.
101. Varghese R, Yesudhasan BL, Vimala LR, Neeravi A, Anandhan K, Baskar P, Elangovan D, Manesh A, James P, Gupta R, et al. Pneumococcal serotypes causing non-invasive pneumonia in adults from a South Indian tertiary care hospital and the impact of the newer conjugate vaccines. *Access Microbiol.* 2021;3(12). doi:10.1099/acmi.0.000258.
102. Oliveira GS, Oliveira MLS, Miyaji EN, Rodrigues TC. Pneumococcal vaccines: past findings, present work, and future strategies. *Vaccines.* 2021;9(11):1338. doi:10.3390/vaccines9111338.
103. Pollard AJ, Perrett KP, Beverley PC. Maintaining protection against invasive bacteria with protein-polysaccharide conjugate vaccines. *Nat Rev Immunol.* 2009;9(3):213–220. doi:10.1038/nri2494.
104. Morais V, Suarez N. Conjugation mechanism for pneumococcal glycoconjugate vaccines: classic and emerging methods. *Bioengineering.* 2022;9(12):774. doi:10.3390/bioengineering9120774.
105. Perera SR, Sokaribo AS, White AP. Polysaccharide vaccines: a perspective on non-typhoidal Salmonella. *Polysaccharides.* 2021;2(3):691–714. doi:10.3390/polysaccharides2030042.
106. PNEUMOVAX 23 dosage and administration. [accessed 2024 Aug 7]. <https://www.merckvaccines.com/vaxneuvancepneumovax23/p23-dosage-and-administration/>.
107. Shapiro ED, Berg AT, Austrian R, Schroeder D, Parcells V, Margolis A, Adair RK, Clemens JD. The protective efficacy of polyvalent pneumococcal polysaccharide vaccine. *N Engl J Med.* 1991;325(21):1453–1460. doi:10.1056/NEJM199111213252101.
108. Jackson LA, Neuzil KM, Yu O, Benson P, Barlow WE, Adams AL, Hanson CA, Mahoney LD, Shay DK, Thompson WW. Effectiveness of pneumococcal polysaccharide vaccine in older adults. *N Engl J Med.* 2003;348(18):1747–1755. doi:10.1056/NEJMoa022678.
109. Maruyama T, Taguchi O, Niederman MS, Morser J, Kobayashi H, Kobayashi T, D'Alessandro-Gabazza C, Nakayama S, Nishikubo K, Noguchi T, et al. Efficacy of 23-valent pneumococcal vaccine in preventing pneumonia and improving survival in nursing home residents: double blind, randomised and placebo controlled trial. *BMJ.* 2010;340(mar08 1):c1004–c1004. doi:10.1136/bmj.c1004.
110. Moberley S, Holden J, Tatham DP, Andrews RM. Vaccines for preventing pneumococcal infection in adults. In: Cochrane Acute Respiratory Infections Group, editor. *The Cochrane Collaboration, ed. Cochrane Database of Systematic Reviews.* Chichester (UK): John Wiley & Sons, Ltd.; 2008. p. CD000422.
111. See KC. Pneumococcal vaccination in adults: a narrative review of considerations for individualized decision-making. *Vaccines.* 2023;11(5):908. doi:10.3390/vaccines11050908.
112. Daniels CC, Rogers PD, Shelton CM. A review of pneumococcal vaccines: current polysaccharide vaccine recommendations and future protein antigens. *J Pediatr Pharmacol Ther.* 2016;21(1):27–35. doi:10.5863/1551-6776-21.1.27.
113. Micoli F, Stefanetti G, MacLennan CA. Exploring the variables influencing the immune response of traditional and innovative glycoconjugate vaccines. *Front Mol Biosci.* 2023;10:1201693. doi:10.3389/fmolb.2023.1201693.
114. Alicino C, Paganino C, Orsi A, Astengo M, Trucchi C, Icardi G, Ansaldi F. The impact of 10-valent and 13-valent pneumococcal conjugate vaccines on hospitalization for pneumonia in children: a systematic review and meta-analysis. *Vaccine.* 2017;35(43):5776–5785. doi:10.1016/j.vaccine.2017.09.005.
115. Reyburn R, Tsatsaronis A, Von Mollendorf C, Mulholland K, Russell FM. ARI Review group. Systematic review on the impact of the pneumococcal conjugate vaccine ten valent (PCV10) or thirteen valent (PCV13) on all-cause, radiologically confirmed and severe pneumonia hospitalisation rates and pneumonia mortality in children 0-9 years old. *J Glob Health.* 2023;13:05002 doi:10.7189/jogh.13.05002.
116. Changes in invasive pneumococcal disease incidence following introduction of PCV10 and 1170 PCV13 in children and adults globally: the pserenade project. [accessed 2024 May 14]. <https://cslide.ctimeetingtech.com/isppd22/attendee/eposter/poster/852>.
117. Weuphoria launching ceremony successfully held. [accessed 2024 May 14]. <https://en.walvax.com/news/weuphoria-launching-ceremony-successfully-held-in-thailand.html>.
118. Tin Tin Htar M, Morato Martínez J, Theilacker C, Schmitt H-J, Swerdlow D. Serotype evolution in Western Europe: perspectives on invasive pneumococcal diseases (IPD). *Expert Rev Vaccines.* 2019;18(11):1145–1155. doi:10.1080/14760584.2019.1688149.
119. Hanquet G, Krizova P, Dalby T, Ladhani SN, Nuorti JP, Danis K, Mereckiene J, Knol MJ, Winje BA, Ciruela P, et al. Serotype replacement after introduction of 10-valent and 13-valent pneumococcal conjugate vaccines in 10 countries, Europe. *Emerg Infect Dis.* 2022;28(1):137–138. doi:10.3201/eid2801.210734.
120. U.S. FDA Approves Merck's VAXNEUVANCE™ (Pneumococcal 15-valent Conjugate Vaccine) for the Prevention of invasive pneumococcal disease in infants and children. Merck.com [accessed 2024 Aug 8]. <https://www.merck.com/news/u-s-fda-approves-mercks-vaxneuvance-pneumococcal-15-valent-conjugate-vaccine-for-the-prevention-of-invasive-pneumococcal-disease-in-infants-and-children/>.
121. Prevnar 20 FDA Approval History. Drugs.com. [accessed 2024 Aug 8]. <https://www.drugs.com/history/prevnar-20.html>.
122. VAXNEUVANCE Prescribing Information. [accessed 2024 May 14]. https://www.merck.com/product/usa/pi_circulars/v/vaxneuvance/vaxneuvance_pi.pdf.
123. Prevnar 20. [accessed 2024 May 14]. <https://www.prevnar20.com/>.
124. U.S. FDA approves PREVNAR 20®, Pfizer's 20-valent Pneumococcal conjugate vaccine for infants and children. [accessed 2024 May 14]. <https://www.pfizer.com/news/press-release/press-release-detail/us-fda-approves-prevnar-20r-pfizers-20-valent-pneumococcal>.

125. U.S. FDA approves PREVNAR 20™, Pfizer's Pneumococcal 20 valent conjugate vaccine for adults ages 18 years or older. [accessed 2024 May 14]. <https://www.pfizer.com/news/press-release/press-release-detail/us-fda-approves-prevnar-20tm-pfizers-pneumococcal-20-valent>.
126. DCGI approves BE's 14-valent pneumococcal conjugate vaccine. [accessed 2024 Aug 8]. <https://www.biologicle.com/news.html>.
127. Tergene Biotech Pvt. Ltd. a Joint Venture company, receives SEC recommendation for pneumococcal 15 valent (PCV15) vaccine. [accessed 2024 Aug 8]. https://nsearchives.nseindia.com/corporate/AUROPHARMA_27122022115948_LtrToSEsPressReleaseTergene27122022.pdf.
128. U.S. FDA approves CAPVAXIVE™ (Pneumococcal 21-valent Conjugate Vaccine) for prevention of invasive pneumococcal disease and pneumococcal pneumonia in adults. Merck.com. [accessed 2024 Aug 8]. <https://www.merck.com/news/u-s-fda-approves-capvaxive-pneumococcal-21-valent-conjugate-vaccine-for-prevention-of-invasive-pneumococcal-disease-and-pneumococcal-pneumonia-in-adults/>.
129. R&D pipeline. [accessed 2024 Aug 8]. <https://www.panaceabiotech.com/en/products/pipeline>.
130. Vaccines. [accessed 2024 Jul 7]. <https://techinvention.biz/rnd-vaccines.html>.
131. SK bioscience-Sanofi announce positive results from phase II study of 21-valent pneumococcal conjugate vaccine candidate. [accessed 2024 Aug 8]. https://www.skbioscience.com/en/news/news_01_01.
132. GSK completes acquisition of Affinivax, Inc. [accessed 2024 May 14]. <https://www.gsk.com/en-gb/media/press-releases/gsk-completes-acquisition-of-affinivax-inc/>.
133. Vaxcyte reports positive data from phase 2 study of its 24-valent pneumococcal conjugate vaccine candidate, VAX-24, in adults aged 65 and older and full six-month safety data from adult phase 1/2 and phase 2 studies. [accessed 2024 May 14]. <https://investors.vaxcyte.com/news-releases/news-release-details/vaxcyte-reports-positive-data-phase-2-study-its-24-valent/>.
134. Vaxcyte doses first participants in phase 1/2 clinical study evaluating VAX-31 for the prevention of invasive pneumococcal disease in adults. [accessed 2024 May 14]. <https://investors.vaxcyte.com/news-releases/news-release-details/vaxcyte-doses-first-participants-phase-12-clinical-study-0/>.
135. Vaxcyte announces FDA clearance of investigational new drug application for VAX-31 for the prevention of invasive pneumococcal disease in adults. [accessed 2024 May 14]. <https://investors.vaxcyte.com/news-releases/news-release-details/vaxcyte-announces-fda-clearance-investigational-new-drug-1/>.
136. McGuinness D, Kaufhold RM, McHugh PM, Winters MA, Smith WJ, Giovarelli C, He J, Zhang Y, Musey L, Skinner JM. Immunogenicity of PCV24, an expanded pneumococcal conjugate vaccine, in adult monkeys and protection in mice. *Vaccine*. 2021;39(30):4231–4237. doi:10.1016/j.vaccine.2021.04.067.
137. Inventprise completes vaccination of participants in a phase 2 dose ranging study of its 25 valent pneumococcal vaccine candidate. [accessed 2024 Aug 9]. <https://inventprise.com/2024/01/02/inventprise-completes-vaccination-of-participants-in-a-phase-2-dose-ranging-study-of-its-25-valent-pneumococcal-vaccine-candidate/>.
138. Conjugate vaccine introduction & use. [accessed 2025 Jan 18]. <https://view-hub.org/vaccine/pcv>.
139. Pneumococcal Conjugate Vaccine (PCV) Review of impact evidence (PRIME) summary of findings from systematic review October 2017. [accessed 2025 Jan 18]. https://terrace.who.int/mediacentre/data/sage/SAGE_Docs_Ppt_Oct2017/9_session_PCV/Oct2019_session9_PCV_PRIMEsummary.pdf.
140. Duke JA, Avci FY. Emerging vaccine strategies against the incessant pneumococcal disease. *NPJ Vaccines*. 2023;8(1):122. doi:10.1038/s41541-023-00715-w.
141. WALVAX Package Insert. [accessed 2024 May 14]. <https://en.walvax.com/media/upload/product/Package%20Insert%20of%20PPSV23.pdf>.
142. Pneumococcal Conjugate Vaccine (PCV). [accessed 2024 May 14]. <https://view-hub.org/vaccine/pcv/product>.
143. Rao MIS, Kasi SG, Dhir SK, Wadhwa A, Rajsekhar B, Kumar CM, Lalwani S, Shenoy B, Kesavan TMA, Kalyani S, et al. Indian Academy of Pediatrics (IAP) Advisory Committee on Vaccines and Immunization Practices (ACVIP): recommended immunization schedule (2023) and update on immunization for children aged 0 through 18 years. *Indian Pediatr*. 2024;61(2):113–125. doi:10.1007/s13312-024-3104-5.
144. Affordable Vaccine platform. [accessed 2024 Aug 8]. <http://www.tergene.com/news3.html>.
145. Curry S, Kaufhold RM, Monslow MA, Zhang Y, McGuinness D, Kim E, Nawrocki DK, McHugh PM, Briggs ML, Smith WJ, et al. Preclinical evaluation of an investigational 21-valent pneumococcal conjugate vaccine, V116, in adult-rhesus monkey, rabbit, and mouse models. *Vaccine*. 2023;41(4):903–913. doi:10.1016/j.vaccine.2022.12.017.
146. Permission to conduct clinical trial of new drug or investigational new drug. [accessed 2024 Aug 8]. https://cdsc.gov.in/opencms/resources/UploadCDSCOWeb/2018/UploadCTApprovals/39_Panacea%20Biotech%20Pneumococcal.pdf.
147. Intellectual property. [accessed 2024 Aug 9]. <https://techinvention.biz/intellectual-property.html>.
148. Composition of multivalent pneumococcal polysaccharide-protein conjugate. [accessed 2024 Aug 8]. <https://pubchem.ncbi.nlm.nih.gov/patent/BR-112020014978-A2>.
149. GSK - A phase 2 study to evaluate the safety, tolerability, and immunogenicity of AFX3772 vaccine in healthy infants. [accessed 2024 May 14]. <https://www.gsk-studyregister.com/en/trial-details/?id=219651>.
150. Fairman J, Agarwal P, Barbanell S, Behrens C, Berges A, Burky J, Davey P, Fernsten P, Grainger C, Guo S, et al. Non-clinical immunological comparison of a Next-Generation 24-valent pneumococcal conjugate vaccine (VAX-24) using site-specific carrier protein conjugation to the current standard of care (PCV13 and PPV23). *Vaccine*. 2021;39(23):3197–3206. doi:10.1016/j.vaccine.2021.03.070.
151. Jain SS, Singh VK, Kante RK, Jana SK, Patil RH. Current trends in development and manufacturing of higher-valent pneumococcal polysaccharide conjugate vaccine and its challenges. *Biol J Int Assoc Biol Stand*. 2024;87:101784. doi:10.1016/j.biologicals.2024.101784.
152. Datta A, Kapre K, Andi-Lolo I, Kapre S. Multi-valent pneumococcal conjugate vaccine for global health: From problem to platform to production. *Hum Vaccines Immunother*. 2022;18(4):2117949. doi:10.1080/21645515.2022.2117949.
153. Behrens C, Fairman J, Agarwal P, Arulkumar S, Barbanell S, Bautista L, Berges A, Burky J, Davey P, Grainger C, et al. 1047. Development of a next generation 30+ valent pneumococcal conjugate vaccine (VAX-XP) using site-specific carrier protein conjugation. *Open Forum Infect Dis*. 2021;8(Supplement_1):S615–S615. doi:10.1093/ofid/ofab466.1241.
154. Pneumococcal Vaccines. [accessed 2024 Aug 28]. <https://www.cdc.gov/vaccines/acip/meetings/downloads/slides-2024-06-26-28/01-Pneumococcal-Loehr-508.pdf>.
155. Feemster K, Buchwald UK, Bannietts N, Joyce JG, Velentgas P, Chapman TJ, Yildirim I. Immunogenicity of current and next generation pneumococcal conjugate vaccines in children: current challenges and upcoming opportunities. *Open Forum Infect Dis*. 2024;11(5):ofae220. doi:10.1093/ofid/ofae220.
156. Wassil J, Sisti M, Fairman J, Rankin B, Clark J, Bennett S, Johnson D, Migone T-S, Nguyen K, Paschenko A, et al. A phase 2, randomized, blinded, dose-finding, controlled clinical trial to evaluate the safety, tolerability, and immunogenicity of a 24-valent pneumococcal conjugate vaccine (VAX-24) in healthy adults 65 years and older. *Vaccine*. 2024;42(25):126124. doi:10.1016/j.vaccine.2024.07.025.
157. Investigational pneumococcal vaccine broadens coverage without reducing immunogenicity. [accessed 2024 Dec 13]. <https://www.contagionlive.com/view/investigational-pneumococcal-vaccine-broadens-coverage-without-reducing-immunogenicity>.

158. Akkoyunlu M. State of pneumococcal vaccine immunity. *Hum Vaccines Immunother.* 2024;20(1):2336358. doi:10.1080/21645515.2024.2336358.
159. Recommendations to assure the quality, safety and efficacy of pneumococcal conjugate vaccines Replacement of WHO Technical Report Series, No. 927, Annex 2. [accessed 2024 Dec 13]. https://cdn.who.int/media/docs/default-source/biologicals/vaccine-standardization/pneumococcus/trs_977_annex_3.pdf.
160. Golden AR, Adam HJ, Karlowsky JA, Baxter M, Nichol KA, Martin I, Demczuk W, Van Caesele P, Gubbay JB, Lefebvre B, et al. Molecular characterization of predominant Streptococcus pneumoniae serotypes causing invasive infections in Canada: the SAVE study, 2011–15. *J Antimicrob Chemother.* 2018;73(suppl_7):vii20–31. doi:10.1093/jac/dky157.
161. Chochua S, Metcalf BJ, Li Z, Walker H, Tran T, McGee L, Beall B. Invasive serotype 35B pneumococci including an expanding serotype switch lineage, United States, 2015–2016. *Emerg Infect Dis.* 2017;23(6):922–930. doi:10.3201/eid2306.170071.
162. Lagousi T, Basdeki P, Routsias J, Spoulou V. Novel protein-based pneumococcal vaccines: assessing the use of distinct protein fragments instead of full-length proteins as vaccine antigens. *Vaccines.* 2019;7(1):9. doi:10.3390/vaccines7010009.
163. Feemster K, Weaver J, Buchwald U, Banniets N, Cox KS, McIntosh ED, Spoulou V. Pneumococcal vaccine breakthrough and failure in infants and children: a narrative review. *Vaccines.* 2023;11(12):1750. doi:10.3390/vaccines11121750.
164. Oligbu G, Hsia Y, Folgori L, Collins S, Ladhani S. Pneumococcal conjugate vaccine failure in children: a systematic review of the literature. *Vaccine.* 2016;34(50):6126–6132. doi:10.1016/j.vaccine.2016.10.050.
165. Aceil J, Avci FY. Pneumococcal surface proteins as virulence factors, immunogens, and conserved vaccine targets. *Front Cell Infect Microbiol.* 2022;12:832254. doi:10.3389/fcimb.2022.832254.
166. Dagan R, Poolman J, Siegrist C-A. Glycoconjugate vaccines and immune interference: a review. *Vaccine.* 2010;28(34):5513–5523. doi:10.1016/j.vaccine.2010.06.026.
167. Musher DM, Anderson R, Feldman C. The remarkable history of pneumococcal vaccination: an ongoing challenge. *Pneumonia.* 2022;14(1):5. doi:10.1186/s41479-022-00097-y.
168. Jang A-Y, Ahn KB, Zhi Y, Ji H-J, Zhang J, Han SH, Guo H, Lim S, Song JY, Lim JH, et al. Serotype-independent protection against invasive pneumococcal infections conferred by live Vaccine with lgt deletion. *Front Immunol.* 2019;10:1212. doi:10.3389/fimmu.2019.01212.
169. Kumraj G, Pathak S, Shah S, Majumder P, Jain J, Bhati D, Hanif S, Mukherjee S, Ahmed S. Capacity building for vaccine manufacturing across developing countries: the way forward. *Hum Vaccines Immunother.* 2022;18(1):2020529. doi:10.1080/21645515.2021.2020529.
170. Nishimoto AT, Rosch JW, Tuomanen EI. Pneumolysin: pathogenesis and therapeutic target. *Front Microbiol.* 2020;11:1543. doi:10.3389/fmicb.2020.01543.
171. Pereira JM, Xu S, Leong JM, Sousa S. The Yin and Yang of pneumolysin during pneumococcal infection. *Front Immunol.* 2022;13:878244. doi:10.3389/fimmu.2022.878244.
172. Araujo AP, Oliveira MLS, Miyaji EN. Negligible role for pneumococcal surface protein a (PspA) and pneumococcal surface protein C (PspC) in the nasopharyngeal colonization of mice with a serotype 6B pneumococcal strain. *Microb Pathog.* 2023;185:106391. doi:10.1016/j.micpath.2023.106391.
173. Du S, Vilhena C, King S, Sahagún-Ruiz A, Hammerschmidt S, Skerka C, Zipfel PF. Molecular analyses identifies new domains and structural differences among Streptococcus pneumoniae immune evasion proteins PspC and Hic. *Sci Rep.* 2021;11(1):1701. doi:10.1038/s41598-020-79362-3.
174. Khan MN, Sharma SK, Filkins LM, Pichichero ME. PcpA of Streptococcus pneumoniae mediates adherence to nasopharyngeal and lung epithelial cells and elicits functional antibodies in humans. *Microbes Infect.* 2012;14(12):1102–1110. doi:10.1016/j.micinf.2012.06.007.
175. Afshar D, Rafiee F, Kheirandish M, Ohadian Moghadam S, Azarsa M. Autolysin (lytA) recombinant protein: a potential target for developing vaccines against pneumococcal infections. *Clin Exp Vaccine Res.* 2020;9(2):76. doi:10.7774/cevr.2020.9.2.76.
176. Hu Y, Park N, Seo KS, Park JY, Somarathne RP, Vonkhalee N, Jackson K, Olivier AK, Fitzkee NC, Thornton JA. Pneumococcal surface adhesion a protein (PsaA) interacts with human Annexin A2 on airway epithelial cells. *Virulence.* 2021;12(1):1947176. doi:10.1080/21505594.2021.1947176.
177. Waz NT, Milani B, Assoni L, Coelho GR, Sciani JM, Parisotto T, Ferraz LFC, Hakansson AP, Converso TR, Darrieux M. Pneumococcal surface protein a (PspA) prevents killing of Streptococcus pneumoniae by indolicidin. *Sci Rep.* 2024;14(1):23517. doi:10.1038/s41598-024-73564-9.
178. Adamou JE, Heinrichs JH, Erwin AL, Walsh W, Gayle T, Dormitzer M, Dagan R, Brewah YA, Barren P, Lathigra R, et al. Identification and characterization of a novel family of pneumococcal proteins that are protective against sepsis. *Infect Immun.* 2001;69(2):949–958. doi:10.1128/IAI.69.2.949-958.2001.
179. Melin M, Di Paolo E, Tikkanen L, Jarva H, Neyt C, Käyhty H, Meri S, Poolman J, Väkeväinen M. Interaction of pneumococcal histidine triad proteins with human complement. *Infect Immun.* 2010;78(5):2089–2098. doi:10.1128/IAI.00811-09.
180. Khan MN, Pichichero ME. Vaccine candidates PhtD and PhtE of Streptococcus pneumoniae are adhesins that elicit functional antibodies in humans. *Vaccine.* 2012;30(18):2900–2907. doi:10.1016/j.vaccine.2012.02.023.
181. Afshari E, Ahangari Cohan R, Shams Nosrati MS, Mousavi SF. Development of a bivalent protein-based vaccine candidate against invasive pneumococcal diseases based on novel pneumococcal surface protein a in combination with pneumococcal histidine triad protein D. *Front Immunol.* 2023;14:1187773. doi:10.3389/fimmu.2023.1187773.
182. Huang J, Gingerich AD, Royer F, Paschall AV, Pena-Briseno A, Avci FY, Mousa JJ. Broadly reactive human monoclonal antibodies targeting the pneumococcal histidine triad protein protect against fatal pneumococcal infection. *Infect Immun.* 2021;89(5):e00747–20. doi:10.1128/IAI.00747-20.
183. Eijkelkamp BA, Pederick VG, Plumtre CD, Harvey RM, Hughes CE, Paton JC, McDevitt CA. The first histidine triad motif of PhtD is critical for Zinc homeostasis in streptococcus pneumoniae. *Infect Immun.* 2016;84(2):407–415. doi:10.1128/IAI.01082-15.
184. Nagai K, Domon H, Maekawa T, Hiyoshi T, Tamura H, Yonezawa D, Habuka R, Saitoh A, Terao Y. Immunization with pneumococcal elongation factor Tu enhances serotype-independent protection against Streptococcus pneumoniae infection. *Vaccine.* 2019;37(1):160–168. doi:10.1016/j.vaccine.2018.11.015.
185. Salha D, Szeto J, Myers L, Claus C, Sheung A, Tang M, Ljutic B, Hanwell D, Ogilvie K, Ming M, et al. Neutralizing antibodies elicited by a novel detoxified pneumolysin derivative, PlyD1, provide protection against both pneumococcal infection and lung injury. *Infect Immun.* 2012;80(6):2212–2220. doi:10.1128/IAI.06348-11.
186. Wu J, Wu K, Xu W, Yuan T, Wang X, Zhang J, Min Y, Yin Y, Zhang X. Engineering detoxified pneumococcal pneumolysin derivative ΔA146PLY for self-biomaterialization of calcium phosphate: assessment of their protective efficacy in murine infection models. *Biomaterials.* 2018;155:152–164. doi:10.1016/j.biomaterials.2017.11.018.
187. Hermand P, Vandercammen A, Mertens E, Di Paolo E, Verlant V, Denoël P, Godfroid F. Preclinical evaluation of a chemically detoxified pneumolysin as pneumococcal vaccine antigen. *Hum Vaccines Immunother.* 2017;13(1):220–228. doi:10.1080/21645515.2016.1234553.
188. Wang Y, Xia L, Wang G, Lu H, Wang H, Luo S, Zhang T, Gao S, Huang J, Min X. Subcutaneous immunization with the fusion

- protein Δ A146Ply-SP0148 confers protection against *Streptococcus pneumoniae* infection. *Microb Pathog.* **2022**;162:105325. doi:10.1016/j.micpath.2021.105325.
189. Kirkham L-A, Kerr AR, Douce GR, Paterson GK, Dilts DA, Liu D-F, Mitchell TJ. Construction and immunological characterization of a novel nontoxic protective pneumolysin mutant for use in future pneumococcal vaccines. *Infect Immun.* **2006**;74(1):586–593. doi:10.1128/IAI.74.1.586-593.2006.
 190. Thanawastien A, Joyce KE, Cartee RT, Haines LA, Pelton SI, Tweten RK, Killeen KP. Preclinical in vitro and in vivo profile of a highly-attenuated, broadly efficacious pneumolysin genetic toxoid. *Vaccine.* **2021**;39(11):1652–1660. doi:10.1016/j.vaccine.2020.04.064.
 191. Verhoeven D, Perry S, Pichichero ME, Plotkin SA. Contributions to protection from streptococcus pneumoniae infection using the monovalent recombinant protein vaccine candidates PcpA, PhtD, and PlyD1 in an infant murine model during challenge. *Clin Vaccine Immunol.* **2014**;21(8):1037–1045. doi:10.1128/CVI.00052-14.
 192. Mann B, Orihuela C, Antikainen J, Gao G, Sublett J, Korhonen TK, Tuomanen E. Multifunctional Role of Choline Binding Protein G in Pneumococcal Pathogenesis. *Infect Immun.* **2006**;74(2):821–829. doi:10.1128/IAI.74.2.821-829.2006.
 193. Kazemian H, Afshar D, Garcia E, Pourmand MR, Jeddi-Tehrani M, Aminharati F, Shokri F, Yazdi MH. CbpM and CbpG of *Streptococcus Pneumoniae* elicit a high protection in Mice challenged with a serotype 19F pneumococcus. *Iran J Allergy Asthma Immunol.* **2018**;17:574–585. doi:10.18502/ijaai.v17i6.622.
 194. Yuan ZQ, Lv ZY, Gan HQ, Xian M, Zhang KX, Mai JY, Yu XB, Wu ZD. Intranasal immunization with autolysin (LytA) in mice model induced protection against five prevalent *Streptococcus pneumoniae* serotypes in China. *Immunol Res.* **2011**;51(1):108. doi:10.1007/s12026-011-8234-x.
 195. Corsini B, Aguinalde L, Ruiz S, Domenech M, Antequera ML, Fenoll A, García P, García E, Yuste J. Immunization with LytB protein of *Streptococcus pneumoniae* activates complement-mediated phagocytosis and induces protection against pneumonia and sepsis. *Vaccine.* **2016**;34(50):6148–6157. doi:10.1016/j.vaccine.2016.11.001.
 196. Balachandran P, Brooks-Walter A, Virolainen-Julkunen A, Hollingshead SK, Briles DE. Role of pneumococcal surface Protein C in Nasopharyngeal carriage and pneumonia and its ability to elicit protection against carriage of *Streptococcus pneumoniae*. *Infect Immun.* **2002**;70(5):2526–2534. doi:10.1128/IAI.70.5.2526-2534.2002.
 197. Ogunniyi AD, Woodrow MC, Poolman JT, Paton JC, Tuomanen EI. Protection against *Streptococcus pneumoniae* elicited by immunization with pneumolysin and CbpA. *Infect Immun.* **2001**;69(10):5997–6003. doi:10.1128/IAI.69.10.5997-6003.2001.
 198. Cao J, Gong Y, Li D, Yin N, Chen T, Xu W, Zhang X, Yin Y. CD4+ T lymphocytes mediated protection against invasive pneumococcal infection induced by mucosal immunization with ClpP and CbpA. *Vaccine.* **2009**;27(21):2838–2844. doi:10.1016/j.vaccine.2009.02.093.
 199. De Lúcia Hernani M, Ferreira PCD, Ferreira DM, Miyaji EN, Ho PL, Oliveira MLS. Nasal immunization of mice with *Lactobacillus casei* expressing the pneumococcal surface protein C primes the immune system and decreases pneumococcal nasopharyngeal colonization in mice. *FEMS Immunol Med Microbiol.* **2011**;62(3):263–272. doi:10.1111/j.1574-695X.2011.00809.x.
 200. Brooks-Walter A, Briles DE, Hollingshead SK, Fischetti VA. The *pspC* gene of *Streptococcus pneumoniae* encodes a polymorphic protein, PspC, which elicits cross-reactive antibodies to PspA 1535 and provides immunity to Pneumococcal Bacteremia. *Infect Immun.* **1999**;67(12):6533–6542. doi:10.1128/IAI.67.12.6533-6542.1999.
 201. Ferreira DM, Darrieux M, Silva DA, Leite LCC, Ferreira JMC, Ho PL, Miyaji EN, Oliveira MLS. Characterization of protective mucosal and systemic immune responses elicited by Pneumococcal surface protein PspA and PspC nasal vaccines against a respiratory pneumococcal challenge in Mice. *Clin Vaccine Immunol.* **2009**;16(5):636–645. doi:10.1128/CVI.00395-08.
 202. Briles DE, Hollingshead SK, King J, Swift A, Braun P, Park M, Ferguson L, Nahm M, Nabors G. Immunization of humans with recombinant Pneumococcal surface protein a (rPspA) elicits antibodies that passively protect Mice from fatal infection with *Streptococcus pneumoniae* bearing heterologous PspA. *J Infect Dis.* **2000**;182(6):1694–1701. doi:10.1086/317602.
 203. Nagano H, Kawabata M, Sugita G, Tsuruhara A, Ohori J, Jimura T, Miyashita K, Kurono Y, Tomonaga K, Briles DE, et al. Transcutaneous immunization with pneumococcal surface protein A in mice. *The Laryngoscope.* **2018**;128(3). doi:10.1002/lary.26971.
 204. Kong IG, Sato A, Yuki Y, Nochi T, Takahashi H, Sawada S, Mejima M, Kurokawa S, Okada K, Sato S, et al. Nanogel-based PspA intranasal vaccine prevents invasive disease and nasal colonization by *Streptococcus pneumoniae*. *Infect Immun.* **2013**;81(5):1625–1634. doi:10.1128/IAI.00240-13.
 205. Figueiredo DBD, Kaneko K, Rodrigues TDC, MacLoughlin R, Miyaji EN, Saleem I, Gonçalves VM. Pneumococcal surface protein A-hybrid nanoparticles protect Mice from lethal challenge after mucosal immunization targeting the lungs. *Pharmaceutics.* **2022**;14(6):1238. doi:10.3390/pharmaceutics14061238.
 206. Wang D, Lu J, Yu J, Hou H, Leenhouts K, Van Roosmalen ML, Gu T, Jiang C, Kong W, Wu Y. A Novel PspA protein vaccine intranasal delivered by bacterium-like particles provides broad protection against pneumococcal pneumonia in Mice. *Immunol Invest.* **2018**;47(4):403–415. doi:10.1080/08820139.2018.1439505.
 207. Lu J, Guo J, Wang D, Yu J, Gu T, Jiang C, Kong W, Wu Y. Broad protective immune responses elicited by bacterium-like particle-based intranasal pneumococcal particle vaccine displaying PspA2 and PspA4 fragments. *Hum Vaccines Immunother.* **2019**;15(2):371–380. doi:10.1080/21645515.2018.1526556.
 208. Anish C, Upadhyay AK, Sehgal D, Panda AK. Influences of process and formulation parameters on powder flow properties and immunogenicity of spray dried polymer particles entrapping recombinant pneumococcal surface protein a. *Int J Pharm.* **2014**;466(1–2):198–210. doi:10.1016/j.jipharm.2014.03.025.
 209. Rodrigues TC, Oliveira MLS, Soares-Schanoski A, Chavez-Rico SL, Figueiredo DB, Gonçalves VM, Ferreira DM, Kunda NK, Saleem IY, Miyaji EN, et al. Mucosal immunization with PspA (Pneumococcal surface protein a)-adsorbed nanoparticles targeting the lungs for protection against pneumococcal infection. *PLoS One.* **2018**;13(1):e0191692. doi:10.1371/journal.pone.0191692.
 210. Tada R, Suzuki H, Takahashi S, Negishi Y, Kiyono H, Kunisawa J, Aramaki Y. Nasal vaccination with pneumococcal surface protein a in combination with cationic liposomes consisting of DOTAP and DC-chol confers antigen-mediated protective immunity against *Streptococcus pneumoniae* infections in mice. *Int Immunopharmacol.* **2018**;61:385–393. doi:10.1016/j.intimp.2018.06.027.
 211. Wagner-Muñiz DA, Haughney SL, Kelly SM, Wannemuehler MJ, Narasimhan B. Room temperature stable PspA-based Nanovaccine induces protective immunity. *Front Immunol.* **2018**;9:325. doi:10.3389/fimmu.2018.00325.
 212. Fukuyama Y, Yuki Y, Katakai Y, Harada N, Takahashi H, Takeda S, Mejima M, Joo S, Kurokawa S, Sawada S, et al. Nanogel-based pneumococcal surface protein a nasal vaccine induces microRNA-associated Th17 cell responses with neutralizing antibodies against *Streptococcus pneumoniae* in macaques. *Mucosal Immunol.* **2015**;8(5):1144–1153. doi:10.1038/mi.2015.5.
 213. Seo S-U, Kim J-J, Yang H, Kwon H-J, Yang J-Y, Curtiss Iii R, Kweon M-N. Effective protection against secondary pneumococcal pneumonia by oral vaccination with attenuated *Salmonella* delivering PspA antigen in mice. *Vaccine.* **2012**;30(48):6816–6823. doi:10.1016/j.vaccine.2012.09.015.
 214. Darrieux M, Miyaji EN, Ferreira DM, Lopes LM, Lopes APY, Ren B, Briles DE, Hollingshead SK, Leite LCC. Fusion proteins containing family 1 and family 2 PspA fragments elicit protection against *Streptococcus pneumoniae* that correlates with antibody-

- mediated enhancement of complement deposition. *Infect Immun.* 2007;75(12):5930–5938. doi:10.1128/IAI.00940-07.
215. Nakahashi-Ouchida R, Uchida Y, Yuki Y, Katakai Y, Yamanoue T, Ogawa H, Muneshige Y, Nakano N, Hanari K, Miyazaki T, et al. A nanogel-based trivalent PspA nasal vaccine protects macaques from intratracheal challenge with pneumococci. *Vaccine.* 2021;39(25):3353–3364. doi:10.1016/j.vaccine.2021.04.069.
 216. Piao Z, Akeda Y, Takeuchi D, Ishii KJ, Ubukata K, Briles DE, Tomono K, Oishi K. Protective properties of a fusion pneumococcal surface protein a (PspA) vaccine against pneumococcal challenge by five different PspA clades in mice. *Vaccine.* 2014;32(43):5607–5613. doi:10.1016/j.vaccine.2014.07.108.
 217. Oliveira MLS, Miyaji EN, Ferreira DM, Moreno AT, Ferreira PCD, Lima FA, Santos FL, Sakauchi MA, Takata CS, Higashi HG, et al. Combination of pneumococcal surface protein a (PspA) with whole cell pertussis vaccine increases protection against Pneumococcal challenge in Mice. *PLOS ONE.* 2010;5(5):e10863. doi:10.1371/journal.pone.0010863.
 218. Talkington DF, Brown BG, Tharpe JA, Koenig A, Russell H. Protection of mice against fatal pneumococcal challenge by immunization with pneumococcal surface adhesin a (PsaA). *Microb Pathog.* 1996;21(1):17–22. doi:10.1006/mpat.1996.0038.
 219. Wang S, Li Y, Shi H, Scarpellini G, Torres-Escobar A, Roland KL, Curtiss R. Immune responses to recombinant pneumococcal PsaA antigen delivered by a live attenuated *Salmonella* vaccine. *Infect Immun.* 2010;78(7):3258–3271. doi:10.1128/IAI.00176-10.
 220. Gor DO, Ding X, Li Q, Sultana D, Mambula SS, Bram RJ, Greenspan NS. Enhanced immunogenicity of pneumococcal surface adhesin a (PsaA) in mice via fusion to recombinant human B lymphocyte stimulator (BLyS). *Biol Direct.* 2011;6(1):9. doi:10.1186/1745-6150-6-9.
 221. Xu J, Dai W, Wang Z, Chen B, Li Z, Fan X. Intranasal vaccination with Chitosan-DNA nanoparticles expressing Pneumococcal surface antigen a protects Mice against nasopharyngeal colonization by *Streptococcus pneumoniae*. *Clin Vaccine Immunol.* 2011;18(1):75–81. doi:10.1128/CVI.00263-10.
 222. Xu J-H, Dai W-J, Chen B, Fan X-Y. Mucosal immunization with PsaA protein, using Chitosan as a delivery system, increases protection against acute Otitis media and invasive infection by *Streptococcus pneumoniae*. *Scand J Immunol.* 2015;81(3):177–185. doi:10.1111/sji.12267.
 223. André GO, Borges MT, Assoni L, Ferraz LFC, Sakshi P, Adamson P, Gordon DL, Ogunniyi AD, Venter H, Converso TR, et al. Protective role of PhtD and its amino and carboxyl fragments against pneumococcal sepsis. *Vaccine.* 2021;39(27):3626–3632. doi:10.1016/j.vaccine.2021.04.068.
 224. Malekan M, Siadat SD, Aghasadeghi M, Shahrokhi N, Afrough P, Behrouzi A, Ahmadi K, Mousavi SF. Evaluation of protective immunity responses against pneumococcal PhtD and its C-terminal in combination with outer-membrane vesicles as adjuvants. *J Med Microbiol.* 2020;69(3):465–477. doi:10.1099/jmm.0.001103.
 225. Khan MN, Pichichero ME, Camilli A. CD4 T Cell Memory and Antibody Responses Directed against the Pneumococcal Histidine Triad Proteins PhtD and PhtE following Nasopharyngeal colonization and immunization and their role in protection against Pneumococcal colonization in Mice. *Infect Immun.* 2013;81(10):3781–3792. doi:10.1128/IAI.00313-13.
 226. Hamel J, Charland N, Pineau I, Ouellet C, Rioux S, Martin D, Brodeur BR. Prevention of Pneumococcal disease in Mice immunized with conserved surface-accessible proteins. *Infect Immun.* 2004;72(5):2659–2670. doi:10.1128/IAI.72.5.2659-2670.2004.
 227. Rose L, Shivshankar P, Hinojosa E, Rodriguez A, Sanchez CJ, Orihuela CJ. Antibodies against PspP, a Novel *Streptococcus pneumoniae* Adhesin, block adhesion and protect Mice against Pneumococcal challenge. *J Infect Dis.* 2008;198(3):375–383. doi:10.1086/589775.
 228. Lee C-J, Lock RA, Andrew PW, Mitchell TJ, Hansman D, Paton JC. Protection of infant mice from challenge with *Streptococcus pneumoniae* type 19F by immunization with a type 19F polysaccharide—pneumolysin conjugate. *Vaccine.* 1994;12(10):875–878. doi:10.1016/0264-410X(94)90028-0.
 229. Michon F, Fusco PC, Minetti CASA, Laude-Sharp M, Uitz C, Huang C-H, D'Ambra AJ, Moore S, Remeta DP, Heron I, et al. Multivalent pneumococcal capsular polysaccharide conjugate vaccines employing genetically detoxified pneumolysin as a carrier protein. *Vaccine.* 1998;16(18):1732–1741. doi:10.1016/S0264-410X(98)00225-4.
 230. Kuo J, Douglas M, Ree HK, Lindberg AA. Characterization of a recombinant pneumolysin and its use as a protein carrier for pneumococcal type 18C conjugate vaccines. *Infect Immun.* 1995;63(7):2706–2713. doi:10.1128/iai.63.7.2706-2713.1995.
 231. Chen Z, Guo R, Xu J, Qiu C. Immunogenicity and protective immunity against otitis media caused by pneumococcus in mice of Hib conjugate vaccine with PsaA protein carrier. *Front Med.* 2016;10(4):490–498. doi:10.1007/s11684-016-0470-y.
 232. Denoël P, Philipp MT, Doyle L, Martin D, Carletti G, Poolman JT. A protein-based pneumococcal vaccine protects rhesus macaques from pneumonia after experimental infection with *Streptococcus pneumoniae*. *Vaccine.* 2011;29(33):5495–5501. doi:10.1016/j.vaccine.2011.05.051.
 233. Hill S, Entwistle C, Pang Y, Joachim M, McIlgorm A, Dalton K, Burbidge P, Colaco C, Brown J, Goldblatt D, et al. Immunogenicity and mechanisms of action of PnuBioVax, a multi-antigen serotype-independent prophylactic vaccine against infection with *Streptococcus pneumoniae*. *Vaccine.* 2018;36(29):4255–4264. doi:10.1016/j.vaccine.2018.05.122.
 234. Chen A, Mann B, Gao G, Heath R, King J, Maissoneuve J, Alderson M, Tate A, Hollingshead SK, Tweten RK, et al. Multivalent Pneumococcal protein vaccines comprising Pneumolysin with epitopes/fragments of CbpA and/or PspA elicit strong and broad protection. *Clin Vaccine Immunol.* 2015;22(10):1079–1089. doi:10.1128/CVI.00293-15.
 235. Briles DE, Ades E, Paton JC, Sampson JS, Carlone GM, Huebner RC, Virolainen A, Swiatlo E, Hollingshead SK. Intranasal immunization of Mice with a mixture of the pneumococcal proteins PsaA and PspA is highly protective against Nasopharyngeal Carriage of *Streptococcus pneumoniae*. *Infect Immun.* 2000;68(2):796–800. doi:10.1128/IAI.68.2.796-800.2000.
 236. Yu J, Chen X, Li B, Gu T, Meng X, Kong W, Wu Y. A pneumococcal vaccine combination with two proteins containing PspA families 1 and 2 can potentially protect against a wide range of *Streptococcus pneumoniae* strains. *Immunol Res.* 2018;66(4):528–536. doi:10.1007/s12026-018-9016-5.
 237. Zane L, Kraschowetz S, Trentini MM, Alves VDS, Araujo SC, Goulart C, Leite LCDC, Gonçalves VM. Peptide linker increased the stability of pneumococcal fusion protein vaccine candidate. *Front Bioeng Biotechnol.* 2023;11:1108300. doi:10.3389/fbioe.2023.1108300.
 238. Mann B, Thornton J, Heath R, Wade KR, Tweten RK, Gao G, El Kasmi K, Jordan JB, Mitrea DM, Kriwacki R, et al. Broadly protective protein-based Pneumococcal vaccine composed of Pneumolysin Toxoid-CbpA peptide recombinant fusion protein. *J Infect Dis.* 2014;209(7):1116–1125. doi:10.1093/infdis/jit502.
 239. Ogunniyi AD, Grabowicz M, Briles DE, Cook J, Paton JC. Development of a vaccine against invasive Pneumococcal disease based on combinations of virulence proteins. *Infect Immun.* 2007;75(1):350–357. doi:10.1128/IAI.01103-06.
 240. Verhoeven D, Xu Q, Pichichero ME. Vaccination with a *Streptococcus pneumoniae* trivalent recombinant PcpA, PhtD and PlyD1 protein vaccine candidate protects against lethal pneumonia in an infant murine model. *Vaccine.* 2014;32(26):3205–3210. doi:10.1016/j.vaccine.2014.04.004.
 241. Xu Q, Surendran N, Verhoeven D, Klapa J, Ochs M, Pichichero ME. Trivalent pneumococcal protein recombinant vaccine protects against lethal *Streptococcus pneumoniae* pneumonia and correlates with phagocytosis by neutrophils during early pathogenesis. *Vaccine.* 2015;33(8):993–1000. doi:10.1016/j.vaccine.2015.01.014.
 242. Xu Q, Pryharski K, Pichichero ME. Trivalent pneumococcal protein vaccine protects against experimental acute otitis media

- caused by *Streptococcus pneumoniae* in an infant murine model. *Vaccine*. 2017;35(2):337–344. doi:10.1016/j.vaccine.2016.11.046.
243. Visan L, Rouleau N, Proust E, Peyrot L, Donadieu A, Ochs M. Antibodies to PcpA and PhtD protect mice against *Streptococcus pneumoniae* by a macrophage- and complement-dependent mechanism. *Hum Vaccines Immunother*. 2018;14(2):489–494. doi:10.1080/21645515.2017.1403698.
 244. Lu J, Sun T, Wang D, Dong Y, Xu M, Hou H, Kong FT, Liang C, Gu T, Chen P, et al. Protective immune responses elicited by fusion protein containing PsaA and PspA fragments. *Immunol Invest*. 2015;44(5):482–496. doi:10.3109/08820139.2015.1037956.
 245. Olafsdottir TA, Lingnau K, Nagy E, Jonsdottir I, Camilli A. Novel protein-based Pneumococcal vaccines administered with the Th1-promoting adjuvant IC31 induce protective immunity against Pneumococcal disease in neonatal. *Infect Immun*. 2012;80(1):461–468. doi:10.1128/IAI.05801-11.
 246. Jomaa M, Yuste J, Paton JC, Jones C, Dougan G, Brown JS. Antibodies to the iron uptake ABC transporter Lipoproteins PiaA and PiuA promote Opsonophagocytosis of *Streptococcus pneumoniae*. *Infect Immun*. 2005;73(10):6852–6859. doi:10.1128/IAI.73.10.6852-6859.2005.
 247. Brown JS, Ogunniyi AD, Woodrow MC, Holden DW, Paton JC, Tuomanen EI. Immunization with components of two iron uptake ABC transporters protects Mice against systemic *Streptococcus pneumoniae* infection. *Infect Immun*. 2001;69(11):6702–6706. doi:10.1128/IAI.69.11.6702-6706.2001.
 248. Jomaa M, Terry S, Hale C, Jones C, Dougan G, Brown J. Immunization with the iron uptake ABC transporter proteins PiaA and PiuA prevents respiratory infection with *Streptococcus pneumoniae*. *Vaccine*. 2006;24(24):5133–5139. doi:10.1016/j.vaccine.2006.04.012.
 249. Malley R, Lipsitch M, Stack A, Saladino R, Fleisher G, Pelton S, Thompson C, Briles D, Anderson P. Intranasal immunization with killed unencapsulated whole cells prevents colonization and invasive disease by capsulated *Pneumococci*. *Infect Immun*. 2001;69(8):4870–4873. doi:10.1128/IAI.69.8.4870-4873.2001.
 250. HogenEsch H, Dunham A, Hansen B, Anderson K, Maisonneuve J-F, Hem SL. Formulation of a killed whole cell pneumococcus vaccine - effect of aluminum adjuvants on the antibody and IL-17 response. *J Immune Based Ther Vaccines*. 2011;9(1):5. doi:10.1186/1476-8518-9-5.
 251. Lu Y-J, Yadav P, Clements JD, Forte S, Srivastava A, Thompson CM, Seid R, Look J, Alderson M, Tate A, et al. Options for inactivation, adjuvant, and route of topical administration of a killed, unencapsulated *Pneumococcal* whole-cell vaccine. *Clin Vaccine Immunol*. 2010;17(6):1005–1012. doi:10.1128/CI.00036-10.
 252. Xu X, Meng J, Wang Y, Zheng J, Wu K, Zhang X, Yin Y, Zhang Q. Serotype-independent protection against pneumococcal infections elicited by intranasal immunization with ethanol-killed pneumococcal strain, SPY1. *J Microbiol*. 2014;52(4):315–323. doi:10.1007/s12275-014-3583-5.
 253. Kim E-H, Choi S-Y, Kwon M-K, Tran TDH, Park S-S, Lee K-J, Bae S-M, Briles DE, Rhee D-K. *Streptococcus pneumoniae* pep27 mutant as a live vaccine for serotype-independent protection in mice. *Vaccine*. 2012;30(11):2008–2019. doi:10.1016/j.vaccine.2011.11.073.
 254. Kim S-J, Seon SH, Luong TT, Ghosh P, Pyo S, Rhee D-K. Immunization with attenuated non-transformable pneumococcal pep27 and comD mutant provides serotype-independent protection against pneumococcal infection. *Vaccine*. 2019;37(1):90–98. doi:10.1016/j.vaccine.2018.11.027.
 255. Amonov M, Simbak N, Wan Hassan W, Ismail SA, Rahman NI, Clarke SC, Yeo CC. Disruption of the cpsE and endA genes attenuates *Streptococcus pneumoniae* virulence: towards the development of a live attenuated vaccine candidate. *Vaccines*. 2020;8(2):187. doi:10.3390/vaccines8020187.
 256. Ramos-Sevillano E, Ercoli G, Guerra-Assunção JA, Felgner P, Ramiro de Assis R, Nakajima R, Goldblatt D, Tetteh KKA, Heyderman RS, Gordon SB, et al. Protective Effect of Nasal Colonisation with Δ cps/piaA and Δ cps/proAbcstreptococcus pneumoniae Strains against Recolonisation and Invasive Infection. *Vaccines*. 2021;9(3):261. doi:10.3390/vaccines9030261.
 257. Kamtchoua T, Bologa M, Hopfer R, Neveu D, Hu B, Sheng X, Corde N, Pouzet C, Zimmermann G, Gurnathan S. Safety and immunogenicity of the pneumococcal pneumolysin derivative PlyD1 in a single-antigen protein vaccine candidate in adults. *Vaccine*. 2013;31(2):327–333. doi:10.1016/j.vaccine.2012.11.005.
 258. Entwisle C, Hill S, Pang Y, Joachim M, McIlgorm A, Colaco C, Goldblatt D, De Gorguet D'Argoeuves P, Bailey C. Safety and immunogenicity of a novel multiple antigen pneumococcal vaccine in adults: A Phase I randomised clinical trial. *Vaccine*. 2017;35(51):7181–7186. doi:10.1016/j.vaccine.2017.10.076.
 259. Prymula R, Pazdiora P, Traskine M, Rüggeberg JU, Borys D. Safety and immunogenicity of an investigational vaccine containing two common pneumococcal proteins in toddlers: a phase II randomized clinical trial. *Vaccine*. 2014;32(25):3025–3034. doi:10.1016/j.vaccine.2014.03.066.
 260. Seiberling M, Bologa M, Brookes R, Ochs M, Go K, Neveu D, Kamtchoua T, Lashley P, Yuan T, Gurnathan S. Safety and immunogenicity of a pneumococcal histidine triad protein D vaccine candidate in adults. *Vaccine*. 2012;30(52):7455–7460. doi:10.1016/j.vaccine.2012.10.080.
 261. Bologa M, Kamtchoua T, Hopfer R, Sheng X, Hicks B, Bixler G, Hou V, Pehlic V, Yuan T, Gurnathan S. Safety and immunogenicity of pneumococcal protein vaccine candidates: monovalent choline-binding protein a (PcpA) vaccine and bivalent PcpA-pneumococcal histidine triad protein D vaccine. *Vaccine*. 2012;30(52):7461–7468. doi:10.1016/j.vaccine.2012.10.076.
 262. Brooks WA, Chang L-J, Sheng X, Hopfer R. Safety and immunogenicity of a trivalent recombinant PcpA, PhtD, and PlyD1 pneumococcal protein vaccine in adults, toddlers, and infants: a phase I randomized controlled study. *Vaccine*. 2015;33(36):4610–4617. doi:10.1016/j.vaccine.2015.06.078.
 263. Pichichero ME. Pneumococcal whole-cell and protein-based vaccines: changing the paradigm. *Expert Rev Vaccines*. 2017;16(12):1181–1190. doi:10.1080/14760584.2017.1393335.
 264. Leroux-Roels I, Devaster J-M, Leroux-Roels G, Verlant V, Henckaerts I, Moris P, Hermand P, Van Belle P, Poolman JT, Vandepapelière P, et al. Adjuvant system AS02V enhances humoral and cellular immune responses to pneumococcal protein PhtD vaccine in healthy young and older adults: Randomised, controlled trials. *Vaccine*. 2015;33(4):577–584. doi:10.1016/j.vaccine.2013.10.052.
 265. Leroux-Roels G, Maes C, De Boever F, Traskine M, Rüggeberg JU, Borys D. Safety, reactogenicity and immunogenicity of a novel pneumococcal protein-based vaccine in adults: a phase I/II randomized clinical study. *Vaccine*. 2014;32(50):6838–6846. doi:10.1016/j.vaccine.2014.02.052.
 266. Pauksens K, Nilsson AC, Caubet M, Pascal TG, Van Belle P, Poolman JT, Vandepapelière PG, Verlant V, Vink PE. Randomized controlled study of the safety and immunogenicity of *Pneumococcal* vaccine formulations containing PhtD and detoxified *Pneumolysin* with alum or adjuvant system AS02V in elderly adults. *Clin Vaccine Immunol*. 2014;21(5):651–660. doi:10.1128/CI.00807-13.
 267. Odotula A, Ota M, Ogundare E, Antonio M, Owiafe P, Worwui A, Greenwood B, Alderson M, Traskine M, Verlant V, et al. Reactogenicity, safety and immunogenicity of a protein-based pneumococcal vaccine in Gambian children aged 2–4 years: a phase II randomized study. *Hum Vaccines Immunother*. 2016;12(2):393–402. doi:10.1080/21645515.2015.1111496.
 268. Odotula A, Ota MOC, Antonio M, Ogundare EO, Saidu Y, Foster-Nyarko E, Owiafe PK, Ceesay F, Worwui A, Idoko OT, et al. Efficacy of a novel, protein-based pneumococcal vaccine against nasopharyngeal carriage of *Streptococcus pneumoniae* in infants: a phase 2, randomized, controlled, observer-blind study. *Vaccine*. 2017;35(19):2531–2542. doi:10.1016/j.vaccine.2017.03.071.
 269. Odotula A, Ota MOC, Antonio M, Ogundare EO, Saidu Y, Owiafe PK, Worwui A, Idoko OT, Owolabi O, Kampmann B,

- et al. Immunogenicity of pneumococcal conjugate vaccine formulations containing pneumococcal proteins, and immunogenicity and reactogenicity of co-administered routine vaccines – a phase II, randomised, observer-blind study in Gambian infants. *Vaccine*. 2019;37(19):2586–2599. doi:10.1016/j.vaccine.2019.03.033.
270. Hammitt LL, Campbell JC, Borys D, Weatherholtz RC, Reid R, Goklish N, Moulton LH, Traskine M, Song Y, Swinnen K, et al. Efficacy, safety and immunogenicity of a pneumococcal protein-based vaccine co-administered with 13-valent pneumococcal conjugate vaccine against acute otitis media in young children: a phase IIb randomized study. *Vaccine*. 2019;37(51):7482–7492. doi:10.1016/j.vaccine.2019.09.076.
 271. Borys D, Rupp R, Smulders R, Chichili GR, Kovanda LL, Santos V, Malinoski F, Siber G, Malley R, Sebastian S. Safety, tolerability and immunogenicity of a novel 24-valent pneumococcal vaccine in toddlers: a phase 1 randomized controlled trial. *Vaccine*. 2024;42(10):2560–2571. doi:10.1016/j.vaccine.2024.02.001.
 272. Chichili GR, Smulders R, Santos V, Cywin B, Kovanda L, Van Sant C, Malinoski F, Sebastian S, Siber G, Malley R. Phase 1/2 study of a novel 24-valent pneumococcal vaccine in healthy adults aged 18 to 64 years and in older adults aged 65 to 85 years. *Vaccine*. 2022;40(31):4190–4198. doi:10.1016/j.vaccine.2022.05.079.
 273. Miyaji EN, Oliveira MLS, Carvalho E, Ho PL. Serotype-independent pneumococcal vaccines. *Cell Mol Life Sci*. 2013;70(18):3303–3326. doi:10.1007/s00018-012-1234-8.
 274. Kim G-L, Seon S-H, Rhee D-K. Pneumonia and Streptococcus pneumoniae vaccine. *Arch Pharm Res*. 2017;40(8):885–893. doi:10.1007/s12272-017-0933-y.
 275. Voß F, Kohler TP, Meyer T, Abdullah MR, van Opzeeland FJ, Saleh M, Michalik S, van Selm S, Schmidt F, de Jonge MI, et al. Intranasal vaccination with lipoproteins confers protection against Pneumococcal colonisation. *Front Immunol*. 2018;9:2405. doi:10.3389/fimmu.2018.02405.
 276. Kohler S, Voß F, Gómez Mejia A, Brown JS, Hammerschmidt S. Pneumococcal lipoproteins involved in bacterial fitness, virulence, and immune evasion. *FEBS Lett*. 2016;590(21):3820–3839. doi:10.1002/1873-3468.12352.
 277. Moffitt K, Malley R. Rationale and prospects for novel pneumococcal vaccines. *Hum Vaccines Immunother*. 2016;12(2):383–392. doi:10.1080/21645515.2015.1087625.
 278. Keech CA, Morrison R, Anderson P, Tate A, Flores J, Goldblatt D, Briles D, Hural J, Malley R, Alderson MR. A Phase 1 Randomized, Placebo-controlled, Observer-blinded trial to evaluate the safety and immunogenicity of inactivated Streptococcus pneumoniae whole-cell vaccine in adults. *Pediatr Infect Dis J*. 2020;39(4):345–351. doi:10.1097/INF.0000000000002567.
 279. Morais V, Texeira E, Suarez N. Next-generation whole-cell Pneumococcal vaccine. *Vaccines*. 2019;7(4):151. doi:10.3390/vaccines7040151.
 280. Dose-Finding Study of S. Pneumoniae Whole Cell Vaccine Adsorbed to Alum (PATH-wSP) in Healthy Kenyan Adults and Toddlers. [accessed 2024 Dec 13]. <https://clinicaltrials.gov/study/NCT02097472>.
 281. A Study of a Vaccine for Pneumonia in Adults and Toddlers in Kenya. [accessed 2024 Dec 13]. <https://clinicaltrials.gov/study/NCT02543892>.
 282. Evaluation of a Booster Dose of Pneumococcal Vaccine Formulations in Young Adults. [accessed 2024 Dec 13]. <https://clinicaltrials.gov/study/NCT00896064>.